

# Foundations of Artificial Intelligence

## 29. Propositional Logic: Basics

Malte Helmert and Gabriele Röger

University of Basel

April 19, 2017

# Classification

classification:

## Propositional Logic

environment:

- **static** vs. dynamic
- **deterministic** vs. non-deterministic vs. stochastic
- **fully** vs. partially vs. not **observable**
- **discrete** vs. continuous
- **single-agent** vs. multi-agent

problem solving method:

- problem-specific vs. **general** vs. learning

(applications also in more complex environments)

# Propositional Logic: Overview

## Chapter overview: propositional logic

- 29. Basics
- 30. Reasoning and Resolution
- 31. DPLL Algorithm
- 32. Local Search and Outlook

# Motivation

# Propositional Logic: Motivation

## propositional logic

- modeling and representing problems and knowledge
- basics for **general** problem descriptions and solving strategies  
( $\rightsquigarrow$  **automated planning**  $\rightsquigarrow$  later in this course)
- allows for automated **reasoning**

**German:** Aussagenlogik, automatisches Schliessen

# Relationship to CSPs

- **previous topic:** constraint satisfaction problems
- satisfiability problem in propositional logic can be viewed as **non-binary CSP** over  $\{\mathbf{F}, \mathbf{T}\}$
- formula encodes constraints
- solution: satisfying assignment of values to variables
- SAT algorithms for this problem:  $\rightsquigarrow$  DPLL (**next week**)

# Propositional Logic: Description of State Spaces

propositional variables for missionaries and cannibals problem:

two-missionaries-on-left-shore

one-cannibal-on-left-shore

boat-on-left-shore

...

- problem description for general problem solvers
- states represented as truth values of atomic **propositions**

**German:** Aussagenvariablen

# Propositional Logic: Intuition

**propositions:** atomic statements over the world  
that cannot be divided further

Propositions with **logical connectives** like  
“and”, “or” and “not” form the propositional formulas.

**German:** logische Verknüpfungen

# Syntax

# Syntax

$\Sigma$  alphabet of propositions

(e.g.,  $\{P, Q, R, \dots\}$  or  $\{X_1, X_2, X_3, \dots\}$ ).

## Definition (propositional formula)

- $\top$  and  $\perp$  are formulas.
- Every proposition in  $\Sigma$  is an (atomic) formula.
- If  $\varphi$  is a formula, then  $\neg\varphi$  is a formula (**negation**).
- If  $\varphi$  and  $\psi$  are formulas, then so are
  - $\varphi \wedge \psi$  (**conjunction**)
  - $\varphi \vee \psi$  (**disjunction**)
  - $\varphi \rightarrow \psi$  (**implication**)

**German:** aussagenlogische Formel, atomare Formel, Konjunktion, Disjunktion, Implikation

**binding strength:**  $(\neg) > (\wedge) > (\vee) > (\rightarrow)$   
(use brackets if necessary)

# Semantics

# Semantics

A formula is **true** or **false**,  
depending on the **interpretation** of the propositions.

## Semantics: Intuition

- A proposition  $p$  is either true or false.  
The truth value of  $p$  is determined by an **interpretation**.
- The truth value of a formula follows from  
the truth values of the propositions.

## Example

$$\varphi = (P \vee Q) \wedge R$$

- If  $P$  and  $Q$  are false, then  $\varphi$  is false  
(independent of the truth value of  $R$ ).
- If  $P$  and  $R$  are true, then  $\varphi$  is true  
(independent of the truth value of  $Q$ ).

# Semantics: Formally

- defined over **interpretation**  $I : \Sigma \rightarrow \{\mathbf{T}, \mathbf{F}\}$
- interpretation  $I$ : **assignment** of propositions in  $\Sigma$
- When is a formula  $\varphi$  true under interpretation  $I$ ?  
symbolically: When does  $I \models \varphi$  hold?

**German:** Interpretation, Belegung

# Semantics: Formally

## Definition ( $I \models \varphi$ )

- $I \models \top$  and  $I \not\models \perp$
- $I \models P$  iff  $I(P) = \mathbf{T}$  for  $P \in \Sigma$
- $I \models \neg\varphi$  iff  $I \not\models \varphi$
- $I \models \varphi \wedge \psi$  iff  $I \models \varphi$  and  $I \models \psi$
- $I \models \varphi \vee \psi$  iff  $I \models \varphi$  or  $I \models \psi$
- $I \models \varphi \rightarrow \psi$  iff  $I \not\models \varphi$  or  $I \models \psi$
- $I \models \Phi$  for a set of formulas  $\Phi$  iff  $I \models \varphi$  for all  $\varphi \in \Phi$

German:  $I$  erfüllt  $\varphi$ ,  $\varphi$  gilt unter  $I$

# Examples

## Example (Interpretation $I$ )

$$I = \{P \mapsto \mathbf{T}, Q \mapsto \mathbf{T}, R \mapsto \mathbf{F}, S \mapsto \mathbf{F}\}$$

## Which formulas are true under $I$ ?

- $\varphi_1 = \neg(P \wedge Q) \wedge (R \wedge \neg S)$ . Does  $I \models \varphi_1$  hold?
- $\varphi_2 = (P \wedge Q) \wedge \neg(R \wedge \neg S)$ . Does  $I \models \varphi_2$  hold?
- $\varphi_3 = (R \rightarrow P)$ . Does  $I \models \varphi_3$  hold?

# Terminology

## Definition (model)

An interpretation  $I$  is called a **model** of  $\varphi$  if  $I \models \varphi$ .

German: Modell

## Definition (satisfiable etc.)

A formula  $\varphi$  is called

- **satisfiable** if there is an interpretation  $I$  such that  $I \models \varphi$ .
- **unsatisfiable** if  $\varphi$  is not satisfiable.
- **falsifiable** if there is an interpretation  $I$  such that  $I \not\models \varphi$ .
- **valid** (= a **tautology**) if  $I \models \varphi$  for all interpretations  $I$ .

German: erfüllbar, unerfüllbar, falsifizierbar,  
allgemeingültig (gültig, Tautologie)

# Terminology

## Definition (logical equivalence)

Formulas  $\varphi$  and  $\psi$  are called **logically equivalent** ( $\varphi \equiv \psi$ ) if for all interpretations  $I$ :  $I \models \varphi$  iff  $I \models \psi$ .

**German:** logisch äquivalent

# Truth Tables

## Truth Tables

How to determine automatically if a given formula is (un)satisfiable, falsifiable, valid?

↪ simple method: **truth tables**

**example:** Is  $\varphi = ((P \vee H) \wedge \neg H) \rightarrow P$  valid?

$P$	$H$	$P \vee H$	$((P \vee H) \wedge \neg H)$	$((P \vee H) \wedge \neg H) \rightarrow P$
F	F	F	F	T
F	T	T	F	T
T	F	T	T	T
T	T	T	F	T

$I \models \varphi$  for all interpretations  $I$   $\rightsquigarrow$   $\varphi$  is valid.

- satisfiability, falsifiability, unsatisfiability?

# Normal Forms

# Normal Forms: Terminology

## Definition (literal)

If  $P \in \Sigma$ , then the formulas  $P$  and  $\neg P$  are called **literals**.

$P$  is called **positive literal**,  $\neg P$  is called **negative literal**.

The **complementary literal** to  $P$  is  $\neg P$  and vice versa.

For a literal  $\ell$ , the complementary literal to  $\ell$  is denoted with  $\bar{\ell}$ .

**German:** Literal, positives/negatives/komplementäres Literal

# Normal Forms: Terminology

## Definition (clause)

A disjunction of 0 or more literals is called a **clause**.

The **empty clause**  $\perp$  is also written as  $\square$ .

Clauses consisting of only one literal are called **unit clauses**.

German: Klausel

## Definition (monomial)

A conjunction of 0 or more literals is called a **monomial**.

German: Monom

# Normal Forms

## Definition (normal forms)

A formula  $\varphi$  is in **conjunctive normal form** (CNF, clause form) if  $\varphi$  is a conjunction of 0 or more clauses:

$$\varphi = \bigwedge_{i=1}^n \left( \bigvee_{j=1}^{m_i} \ell_{i,j} \right)$$

A formula  $\varphi$  is in **disjunctive normal form** (DNF) if  $\varphi$  is a disjunction of 0 or more monomials:

$$\varphi = \bigvee_{i=1}^n \left( \bigwedge_{j=1}^{m_i} \ell_{i,j} \right)$$

**German:** konjunktive Normalform, disjunktive Normalform

# Normal Forms

For every propositional formula, there exists a logically equivalent propositional formula in CNF and in DNF.

## Conversion to CNF

important rules for conversion to CNF:

- $\varphi \rightarrow \psi \equiv \neg\varphi \vee \psi$  (( $\rightarrow$ )-elimination)
- $\neg(\varphi \wedge \psi) \equiv \neg\varphi \vee \neg\psi$  (De Morgan)
- $\neg(\varphi \vee \psi) \equiv \neg\varphi \wedge \neg\psi$  (De Morgan)
- $\neg\neg\varphi \equiv \varphi$  (double negation)
- $(\varphi \wedge \psi) \vee \eta \equiv (\varphi \vee \eta) \wedge (\psi \vee \eta)$  (distributivity)

There are formulas  $\varphi$  for which every logically equivalent formula in CNF and DNF is exponentially longer than  $\varphi$ .

# Summary

# Summary (1)

- **Propositional logic** forms the basis for a general representation of problems and knowledge.
- **Propositions** (atomic formulas) are statements over the world which cannot be divided further.
- **Propositional formulas** combine atomic formulas with  $\neg$ ,  $\wedge$ ,  $\vee$ ,  $\rightarrow$  to more complex statements.
- **Interpretations** determine which atomic formulas are true and which ones are false.

## Summary (2)

- important terminology:
  - model
  - satisfiable, unsatisfiable, falsifiable, valid
  - logically equivalent
- different kinds of formulas:
  - atomic formulas and literals
  - clauses and monomials
  - conjunctive normal form and disjunctive normal form