Foundations of Artificial Intelligence 22. Constraint Satisfaction Problems: Introduction and Examples

Malte Helmert

University of Basel

April 12, 2021

Classification

Classification:

Constraint Satisfaction Problems

environment:

- static vs. dynamic
- deterministic vs. non-deterministic vs. stochastic
- fully vs. partially vs. not observable
- discrete vs. continuous
- single-agent vs. multi-agent

problem solving method:

• problem-specific vs. general vs. learning

Constraint Satisfaction Problems: Overview

Chapter overview: constraint satisfaction problems

- 22.-23. Introduction
 - 22. Introduction and Examples
 - 23. Constraint Networks
- 24.-26. Basic Algorithms
- 27.-28. Problem Structure

Introduction

Constraints

What is a Constraint?

a condition that every solution to a problem must satisfy

German: Einschränkung, Nebenbedingung (math.)

Examples: Where do constraints occur?

- mathematics: requirements on solutions of optimization problems (e.g., equations, inequalities)
- software testing: specification of invariants to check data consistency (e.g., assertions)
- databases: integrity constraints

Constraint Satisfaction Problems: Informally

Given:

- set of variables with corresponding domains
- set of constraints that the variables must satisfy
 - most commonly binary, i.e., every constraint refers to two variables

Solution:

• assignment to the variables that satisfies all constraints

German: Variablen, Constraints, binär, Belegung

Examples

Examples

Examples

- 8 queens problem
- Latin squares
- Sudoku
- graph coloring
- satisfiability in propositional logic

German: 8-Damen-Problem, lateinische Quadrate, Sudoku, Graphfärbung, Erfüllbarkeitsproblem der Aussagenlogik

more complex examples:

- systems of equations and inequalities
- database queries

Example: 8 Queens Problem (Reminder)

(reminder from previous two chapters)

8 Queens Problem

How can we

- place 8 queens on a chess board
- such that no two queens threaten each other?
- originally proposed in 1848
- variants: board size; other pieces; higher dimension

There are 92 solutions, or 12 solutions if we do not count symmetric solutions (under rotation or reflection) as distinct.

Examples 0000000000

Summary 00

8 Queens Problem: Example Solution



example solution for the 8 queens problem

Example: Latin Squares

Latin Squares

How can we

- build an n × n matrix with n symbols
- such that every symbol occurs exactly once in every row and every column?

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \\ 3 & 4 & 1 & 2 \\ 4 & 1 & 2 & 3 \end{bmatrix}$$

There exist 12 different Latin squares of size 3, 576 of size 4, 161 280 of size 5, ..., 5 524 751 496 156 892 842 531 225 600 of size 9.

Example: Sudoku

Sudoku

How can we

- completely fill an already partially filled 9 \times 9 matrix with numbers between 1–9
- such that each row, each column, and each of the nine 3×3 blocks contains every number exactly once?

2	5			3		9		1
	1				4			
4		7				2		8
		5	2					
				9	8	1		
	4				3			
			3	6			7	2
	7							3
9		3				6		4

Example: Sudoku

Sudoku

How can we

- completely fill an already partially filled 9 \times 9 matrix with numbers between 1–9
- such that each row, each column, and each of the nine 3×3 blocks contains every number exactly once?



Example: Sudoku

Sudoku

How can we

- completely fill an already partially filled 9 \times 9 matrix with numbers between 1–9
- such that each row, each column, and each of the nine 3×3 blocks contains every number exactly once?



relationship to Latin squares?

Sudoku: Trivia

- well-formed Sudokus have exactly one solution
- to achieve well-formedness, \geq 17 cells must be filled already (McGuire et al., 2012)
- 6 670 903 752 021 072 936 960 solutions
- only 5472730538 "non-symmetrical" solutions

Example: Graph Coloring

Graph Coloring

How can we

- color the vertices of a given graph using k colors
- such that two neighboring vertices never have the same color?

(The graph and k are problem parameters.)

Example: Graph Coloring

Graph Coloring

How can we

- color the vertices of a given graph using k colors
- such that two neighboring vertices never have the same color?

(The graph and k are problem parameters.)

NP-complete problem

- even for the special case of planar graphs and k = 3
- easy for k = 2 (also for general graphs)

Example: Graph Coloring

Graph Coloring

How can we

- color the vertices of a given graph using k colors
- such that two neighboring vertices never have the same color?

(The graph and k are problem parameters.)

NP-complete problem

- even for the special case of planar graphs and k = 3
- easy for k = 2 (also for general graphs)

Relationship to Sudoku?

Four Color Problem

famous problem in mathematics: Four Color Problem

- Is it always possible to color a planar graph with 4 colors?
- conjectured by Francis Guthrie (1852)
- 1890 first proof that 5 colors suffice
- several wrong proofs surviving for over 10 years

Summary 00

Four Color Problem

famous problem in mathematics: Four Color Problem

- Is it always possible to color a planar graph with 4 colors?
- conjectured by Francis Guthrie (1852)
- 1890 first proof that 5 colors suffice
- several wrong proofs surviving for over 10 years
- solved by Appel and Haken in 1976: 4 colors suffice
- Appel and Haken reduced the problem to 1936 cases, which were then checked by computers
- first famous mathematical problem solved (partially) by computers

 \rightsquigarrow led to controversy: is this a mathematical proof?

Numberphile video:

https://www.youtube.com/watch?v=NgbK43jB4rQ

Satisfiability in Propositional Logic

Satisfiability in Propositional Logic

How can we

- assign truth values (true/false) to a set of propositional variables
- such that a given set of clauses
 (formulas of the form X ∨ ¬Y ∨ Z) is satisfied (true)?

Satisfiability in Propositional Logic

Satisfiability in Propositional Logic

How can we

- assign truth values (true/false) to a set of propositional variables
- such that a given set of clauses
 (formulas of the form X ∨ ¬Y ∨ Z) is satisfied (true)?

remarks:

- NP-complete (Cook 1971; Levin 1973)
- formulas expressed as clauses (instead of arbitrary propositional formulas) is no restriction
- clause length bounded by 3 would not be a restriction

Satisfiability in Propositional Logic

Satisfiability in Propositional Logic

How can we

- assign truth values (true/false) to a set of propositional variables
- such that a given set of clauses
 (formulas of the form X ∨ ¬Y ∨ Z) is satisfied (true)?

remarks:

- NP-complete (Cook 1971; Levin 1973)
- formulas expressed as clauses (instead of arbitrary propositional formulas) is no restriction
- ${\ensuremath{\, \bullet }}$ clause length bounded by 3 would not be a restriction

relationship to previous problems (e.g., Sudoku)?

Practical Applications

- There are thousands of practical applications of constraint satisfaction problems.
- This statement is true already for the satisfiability problem of propositional logic.

some examples:

- verification of hardware and software
- timetabling (e.g., generating time schedules, room assignments for university courses)
- assignment of frequency spectra (e.g., broadcasting, mobile phones)

Summary

Summary

o constraint satisfaction:

- find assignment for a set of variables
- with given variable domains
- that satisfies a given set of constraints.

• examples:

- 8 queens problem
- Latin squares
- Sudoku
- graph coloring
- satisfiability in propositional logic
- many practical applications