

# Theory of Computer Science


## C6. Rice's Theorem

Gabriele Röger

University of Basel

April 24, 2024

# Content of the Course



# Rice's Theorem

# Rice's Theorem (1)

- We have shown that the following problems are undecidable:
  - halting problem  $H$
  - halting problem on empty tape  $H_0$
  - post correspondence problem PCP
- Many more results of this type could be shown.
- Instead, we prove a much more general result, **Rice's theorem**, which shows that a very large class of different problems are undecidable.
- Rice's theorem can be summarized informally as:  
**every** non-trivial question about **what** a given Turing machine computes is undecidable.

## Rice's Theorem (2)

### Theorem (Rice's Theorem)

Let  $\mathcal{R}$  be the class of all computable partial functions.

Let  $S$  be an *arbitrary* subset of  $\mathcal{R}$  except  $S = \emptyset$  or  $S = \mathcal{R}$ .

Then the language

$$C(S) = \{w \in \{0, 1\}^* \mid \text{the (partial) function computed by } M_w \\ \text{is in } S\}$$

is undecidable.

**Question:** why the restriction to  $S \neq \emptyset$  and  $S \neq \mathcal{R}$ ?

**Extension (without proof):** in most cases neither  $C(S)$  nor  $\overline{C(S)}$  is Turing-recognizable. (But there are sets  $S$  for which one of the two languages is Turing-recognizable.)

## Rice's Theorem (3)

Proof.

Let  $\Omega$  be the partial function that is undefined everywhere.

## Rice's Theorem (3)

### Proof.

Let  $\Omega$  be the partial function that is undefined everywhere.

Case distinction:

Case 1:  $\Omega \in \mathcal{S}$

# Rice's Theorem (3)

## Proof.

Let  $\Omega$  be the partial function that is undefined everywhere.

Case distinction:

Case 1:  $\Omega \in \mathcal{S}$

Let  $q \in \mathcal{R} \setminus \mathcal{S}$  be an arbitrary computable partial function outside of  $\mathcal{S}$  (exists because  $\mathcal{S} \subseteq \mathcal{R}$  and  $\mathcal{S} \neq \mathcal{R}$ ).



## Rice's Theorem (3)

### Proof.

Let  $\Omega$  be the partial function that is undefined everywhere.

Case distinction:

Case 1:  $\Omega \in \mathcal{S}$

Let  $q \in \mathcal{R} \setminus \mathcal{S}$  be an arbitrary computable partial function outside of  $\mathcal{S}$  (exists because  $\mathcal{S} \subseteq \mathcal{R}$  and  $\mathcal{S} \neq \mathcal{R}$ ).

Let  $Q$  be a Turing machine that computes  $q$ .

...

## Rice's Theorem (4)

### Proof (continued).

We show that  $\bar{H}_0 \leq C(S)$ .

Consider function  $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ ,

where  $f(w)$  is defined as follows:

- Construct TM  $M$  that first behaves on input  $y$  like  $M_w$  on the empty tape (independently of what  $y$  is).
- Afterwards (if that computation terminates!)  $M$  clears the tape, creates the start configuration of  $Q$  for input  $y$  and then simulates  $Q$ .
- $f(w)$  is the encoding of this TM  $M$

## Rice's Theorem (4)

### Proof (continued).

We show that  $\bar{H}_0 \leq C(S)$ .

Consider function  $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ ,

where  $f(w)$  is defined as follows:

- Construct TM  $M$  that first behaves on input  $y$  like  $M_w$  on the empty tape (independently of what  $y$  is).
- Afterwards (if that computation terminates!)  $M$  clears the tape, creates the start configuration of  $Q$  for input  $y$  and then simulates  $Q$ .
- $f(w)$  is the encoding of this TM  $M$

$f$  is total and computable.

...

## Rice's Theorem (5)

Proof (continued).

Which function is computed by the TM encoded by  $f(w)$ ?

# Rice's Theorem (5)

Proof (continued).

Which function is computed by the TM encoded by  $f(w)$ ?

$M_{f(w)}$  computes  $\begin{cases} \Omega & \text{if } M_w \text{ does not terminate on } \varepsilon \\ q & \text{otherwise} \end{cases}$

# Rice's Theorem (5)

Proof (continued).

Which function is computed by the TM encoded by  $f(w)$ ?

$M_{f(w)}$  computes  $\begin{cases} \Omega & \text{if } M_w \text{ does not terminate on } \varepsilon \\ q & \text{otherwise} \end{cases}$

For all words  $w \in \{0, 1\}^*$ :

$w \in H_0 \implies M_w$  terminates on  $\varepsilon$

$\implies M_{f(w)}$  computes the function  $q$

$\implies$  the function computed by  $M_{f(w)}$  is not in  $\mathcal{S}$

$\implies f(w) \notin C(\mathcal{S})$

...

# Rice's Theorem (6)

Proof (continued).

Further:

- $w \notin H_0 \implies M_w$  does not terminate on  $\varepsilon$
- $\implies M_{f(w)}$  computes the function  $\Omega$
- $\implies$  the function computed by  $M_{f(w)}$  is in  $\mathcal{S}$
- $\implies f(w) \in C(\mathcal{S})$

# Rice's Theorem (6)

Proof (continued).

Further:

- $w \notin H_0 \implies M_w$  does not terminate on  $\varepsilon$
- $\implies M_{f(w)}$  computes the function  $\Omega$
- $\implies$  the function computed by  $M_{f(w)}$  is in  $\mathcal{S}$
- $\implies f(w) \in C(\mathcal{S})$

Together this means:  $w \notin H_0$  iff  $f(w) \in C(\mathcal{S})$ ,  
thus  $w \in \bar{H}_0$  iff  $f(w) \in C(\mathcal{S})$ .



# Rice's Theorem (6)

Proof (continued).

Further:

- $w \notin H_0 \implies M_w$  does not terminate on  $\varepsilon$
- $\implies M_{f(w)}$  computes the function  $\Omega$
- $\implies$  the function computed by  $M_{f(w)}$  is in  $\mathcal{S}$
- $\implies f(w) \in C(\mathcal{S})$

Together this means:  $w \notin H_0$  iff  $f(w) \in C(\mathcal{S})$ ,  
thus  $w \in \bar{H}_0$  iff  $f(w) \in C(\mathcal{S})$ .

Therefore,  $f$  is a reduction of  $\bar{H}_0$  to  $C(\mathcal{S})$ .

# Rice's Theorem (6)

Proof (continued).

Further:

- $w \notin H_0 \implies M_w$  does not terminate on  $\varepsilon$
- $\implies M_{f(w)}$  computes the function  $\Omega$
- $\implies$  the function computed by  $M_{f(w)}$  is in  $\mathcal{S}$
- $\implies f(w) \in C(\mathcal{S})$

Together this means:  $w \notin H_0$  iff  $f(w) \in C(\mathcal{S})$ ,  
thus  $w \in \bar{H}_0$  iff  $f(w) \in C(\mathcal{S})$ .

Therefore,  $f$  is a reduction of  $\bar{H}_0$  to  $C(\mathcal{S})$ .

Since  $H_0$  is undecidable,  $\bar{H}_0$  is also undecidable.

# Rice's Theorem (6)

Proof (continued).

Further:

- $w \notin H_0 \implies M_w$  does not terminate on  $\varepsilon$
- $\implies M_{f(w)}$  computes the function  $\Omega$
- $\implies$  the function computed by  $M_{f(w)}$  is in  $\mathcal{S}$
- $\implies f(w) \in C(\mathcal{S})$

Together this means:  $w \notin H_0$  iff  $f(w) \in C(\mathcal{S})$ ,  
thus  $w \in \bar{H}_0$  iff  $f(w) \in C(\mathcal{S})$ .

Therefore,  $f$  is a reduction of  $\bar{H}_0$  to  $C(\mathcal{S})$ .

Since  $H_0$  is undecidable,  $\bar{H}_0$  is also undecidable.

We can conclude that  $C(\mathcal{S})$  is undecidable.

...

# Rice's Theorem (7)

Proof (continued).

Case 2:  $\Omega \notin \mathcal{S}$

Analogous to [Case 1](#) but this time choose  $q \in \mathcal{S}$ .

The corresponding function  $f$  then reduces  $H_0$  to  $C(\mathcal{S})$ .

Thus, it also follows in this case that  $C(\mathcal{S})$  is undecidable. □

# Rice's Theorem: Consequences

## Was it worth it?

We can now conclude immediately that (for example) the following informally specified problems are all undecidable:

- Does a given TM compute a constant function?
- Does a given TM compute a total function (i. e. will it always terminate, and in particular terminate in a “correct” configuration)?
- Is the output of a given TM always longer than its input?
- Does a given TM compute the identity function?
- Does a given TM compute the computable function  $f$ ?
- ...

# Rice's Theorem: Examples

- Does a given TM compute a constant function?

$$\mathcal{S} = \{f \mid f \text{ is total and computable and} \\ \text{for all } x, y \text{ in the domain of } f : f(x) = f(y)\}$$

# Rice's Theorem: Examples

- Does a given TM compute a constant function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable and}$   
for all  $x, y$  in the domain of  $f : f(x) = f(y)\}$
- Does a given TM compute a total function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable}\}$

# Rice's Theorem: Examples

- Does a given TM compute a constant function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable and}$   
for all  $x, y$  in the domain of  $f : f(x) = f(y)\}$
- Does a given TM compute a total function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable}\}$
- Does a given TM compute the identity function?  
 $\mathcal{S} = \{f \mid f(x) = x \text{ for all } x\}$



# Rice's Theorem: Examples

- Does a given TM compute a constant function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable and}$   
for all  $x, y$  in the domain of  $f : f(x) = f(y)\}$
- Does a given TM compute a total function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable}\}$
- Does a given TM compute the identity function?  
 $\mathcal{S} = \{f \mid f(x) = x \text{ for all } x\}$
- Does a given TM add two natural numbers?  
 $\mathcal{S} = \{f : \mathbb{N}_0^2 \rightarrow \mathbb{N}_0 \mid f(x, y) = x + y\}$

# Rice's Theorem: Examples

- Does a given TM compute a constant function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable and}$   
for all  $x, y$  in the domain of  $f : f(x) = f(y)\}$
- Does a given TM compute a total function?  
 $\mathcal{S} = \{f \mid f \text{ is total and computable}\}$
- Does a given TM compute the identity function?  
 $\mathcal{S} = \{f \mid f(x) = x \text{ for all } x\}$
- Does a given TM add two natural numbers?  
 $\mathcal{S} = \{f : \mathbb{N}_0^2 \rightarrow \mathbb{N}_0 \mid f(x, y) = x + y\}$
- Does a given TM compute the computable function  $f$ ?  
 $\mathcal{S} = \{f\}$   
(full automization of software verification is impossible)

# Rice's Theorem: Pitfalls

- $\mathcal{S} = \{f \mid f \text{ can be computed by a DTM with an even number of states}\}$   
Rice's theorem not applicable because  $\mathcal{S} = \mathcal{R}$

# Rice's Theorem: Pitfalls

- $\mathcal{S} = \{f \mid f \text{ can be computed by a DTM with an even number of states}\}$   
Rice's theorem not applicable because  $\mathcal{S} = \mathcal{R}$
- $\mathcal{S} = \{f : \{0,1\}^* \rightarrow_p \{0,1\} \mid f(w) = 1 \text{ iff } M_w \text{ does not terminate on } \epsilon\}$ ?  
Rice's theorem not applicable because  $\mathcal{S} \not\subseteq \mathcal{R}$

# Rice's Theorem: Pitfalls


- $\mathcal{S} = \{f \mid f \text{ can be computed by a DTM with an even number of states}\}$   
Rice's theorem not applicable because  $\mathcal{S} = \mathcal{R}$
- $\mathcal{S} = \{f : \{0,1\}^* \rightarrow_p \{0,1\} \mid f(w) = 1 \text{ iff } M_w \text{ does not terminate on } \epsilon\}$   
Rice's theorem not applicable because  $\mathcal{S} \not\subseteq \mathcal{R}$
- Show that  $\{w \mid M_w \text{ traverses all states on every input}\}$  is undecidable.  
Rice's theorem not directly applicable because not a semantic property (the function computed by  $M_w$  can also be computed by a TM that does not traverse all states)

# Rice's Theorem: Practical Applications

Undecidable due to Rice's theorem + a small reduction:

- **automated debugging:**
  - Can a given variable ever receive a `null` value?
  - Can a given assertion in a program ever trigger?
  - Can a given buffer ever overflow?
- **virus scanners and other software security analysis:**
  - Can this code do something harmful?
  - Is this program vulnerable to SQL injections?
  - Can this program lead to a privilege escalation?
- **optimizing compilers:**
  - Is this dead code?
  - Is this a constant expression?
  - Can pointer aliasing happen here?
  - Is it safe to parallelize this code path?
- **parallel program analysis:**
  - Is a deadlock possible here?
  - Can a race condition happen here?

# Questions



Questions?

# Further Undecidable Problems



## And What Else?

- Here we conclude our discussion of undecidable problems.
- Many more undecidable problems exist.
- In this section, we briefly discuss some further classical results.

# Undecidable Grammar Problems

## Some Grammar Problems

Given context-free grammars  $G_1$  and  $G_2, \dots$

- ... is  $\mathcal{L}(G_1) \cap \mathcal{L}(G_2) = \emptyset$ ?
- ... is  $|\mathcal{L}(G_1) \cap \mathcal{L}(G_2)| = \infty$ ?
- ... is  $\mathcal{L}(G_1) \cap \mathcal{L}(G_2)$  context-free?
- ... is  $\mathcal{L}(G_1) \subseteq \mathcal{L}(G_2)$ ?
- ... is  $\mathcal{L}(G_1) = \mathcal{L}(G_2)$ ?

Given a context-sensitive grammar  $G, \dots$

- ... is  $\mathcal{L}(G) = \emptyset$ ?
- ... is  $|\mathcal{L}(G)| = \infty$ ?

$\rightsquigarrow$  all undecidable by reduction from PCP  
(see Schöning, Chapter 2.8)

# Gödel's First Incompleteness Theorem (1)

## Definition (Arithmetic Formula)

An **arithmetic formula** is a closed predicate logic formula using

- constant symbols 0 and 1,
- function symbols + and  $\cdot$ , and
- equality (=) as the only relation symbols.

It is called **true** if it is true under the usual interpretation of 0, 1, + and  $\cdot$  over  $\mathbb{N}_0$ .

**Example:**  $\forall x \exists y \forall z (((x \cdot y) = z) \wedge ((1 + x) = (x \cdot y)))$

## Gödel's First Incompleteness Theorem (2)


### Gödel's First Incompleteness Theorem

The problem of **deciding if a given arithmetic formula is true** is undecidable.

Moreover, neither it nor its complement are Turing-recognizable.

As a consequence, there exists no sound and complete proof system for arithmetic formulas.

# Questions



Questions?

# Summary

# Summary

## Rice's theorem:

- “In general one cannot determine algorithmically what a given program (or Turing machine) computes.”

# Summary

## Rice's theorem:

- “In general one cannot determine algorithmically what a given program (or Turing machine) computes.”

## How to Prove Undecidability?

- statements on the computed function of a TM/an algorithm
- other problems



# Summary

## Rice's theorem:

- “In general one cannot determine algorithmically what a given program (or Turing machine) computes.”

## How to Prove Undecidability?

- statements on the computed function of a TM/an algorithm  
→ easiest with [Rice' theorem](#)
- other problems

# Summary

## Rice's theorem:

- “In general one cannot determine algorithmically what a given program (or Turing machine) computes.”

## How to Prove Undecidability?

- statements on the computed function of a TM/an algorithm  
→ easiest with [Rice' theorem](#)
- other problems
  - [directly with the definition of undecidability](#)  
→ usually quite complicated

# Summary

## Rice's theorem:

- “In general one cannot determine algorithmically what a given program (or Turing machine) computes.”

## How to Prove Undecidability?

- statements on the computed function of a TM/an algorithm  
→ easiest with [Rice' theorem](#)
- other problems
  - [directly with the definition of undecidability](#)  
→ usually quite complicated
  - [reduction from an undecidable problem](#), e.g.  
→ halting problem ( $H$ )  
→ Post correspondence problem (PCP)

# What's Next?

contents of this course:

- A. **background** ✓
  - ▷ mathematical foundations and proof techniques
- B. **automata theory and formal languages** ✓
  - ▷ What is a computation?
- C. **Turing computability**
  - ▷ What can be computed at all?
- D. **complexity theory**
  - ▷ What can be computed efficiently?
- E. **more computability theory**
  - ▷ Other models of computability

# What's Next?

contents of this course:

- A. **background** ✓
  - ▷ mathematical foundations and proof techniques
- B. **automata theory and formal languages** ✓
  - ▷ What is a computation?
- C. **Turing computability** ✓
  - ▷ What can be computed at all?
- D. **complexity theory**
  - ▷ What can be computed efficiently?
- E. **more computability theory**
  - ▷ Other models of computability

# Quiz



kahoot.it