

# Theory of Computer Science

## A3. Proof Techniques

Gabriele Röger

University of Basel

February 24, 2025

# Theory of Computer Science

February 24, 2025 — A3. Proof Techniques

## A3.1 Proofs & Proof Strategies

## A3.2 Direct Proof

## A3.3 Indirect Proof

## A3.4 Structural Induction

## A3.5 Summary

# A3.1 Proofs & Proof Strategies

# What is a Proof?

A **mathematical proof** is

- ▶ a sequence of logical steps
- ▶ starting with one set of statements
- ▶ that comes to the conclusion  
that some statement must be true.

What is a **statement**?

# Mathematical Statements

## Mathematical Statement

A **mathematical statement** consists of a set of **preconditions** and a set of **conclusions**.

The statement is **true** if the conclusions are true whenever the preconditions are true.

The set of preconditions is sometimes empty.

German: Mathematische Aussage

# Examples of Mathematical Statements

Examples (some true, some false):

- ▶ “Let  $p \in \mathbb{N}_0$  be a prime number. Then  $p$  is odd.”
- ▶ “There exists an even prime number.”
- ▶ “Let  $p \in \mathbb{N}_0$  be a prime number with  $p \geq 3$ . Then  $p$  is odd.”
- ▶ “All prime numbers  $p \geq 3$  are odd.”
- ▶ “If 4 is a prime number then  $2 \cdot 3 = 4$ .”

What are the preconditions, what are the conclusions?

# On what Statements can we Build the Proof?

A mathematical proof is

- ▶ a sequence of logical steps
- ▶ **starting with one set of statements**
- ▶ that comes to the conclusion  
that some statement must be true.

We can use:

- ▶ **axioms**: statements that are assumed to always be true  
in the current context
- ▶ **theorems** and **lemmas**: statements that were already proven
  - ▶ lemma: an intermediate tool
  - ▶ theorem: itself a relevant result
- ▶ **premises**: assumptions we make  
to see what consequences they have

German: Axiom, Theorem/Satz, Lemma, Prämisse/Annahme

# What is a Logical Step?

A mathematical proof is

- ▶ a sequence of logical steps
- ▶ starting with one set of statements
- ▶ that comes to the conclusion that some statement must be true.

Each step **directly follows**

- ▶ from the axioms,
- ▶ premises,
- ▶ previously proven statements and
- ▶ the preconditions of the statement we want to prove.

For a formal definition, we would need formal logics.



# The Role of Definitions

## Definition

A **set** is an unordered collection of distinct objects.

The objects in a set are called the **elements** of the set. A set is said to **contain** its elements.

We write  $x \in S$  to indicate that  $x$  is an element of set  $S$ , and  $x \notin S$  to indicate that  $S$  does not contain  $x$ .

The set that does not contain any objects is the **empty set**  $\emptyset$ .

- ▶ A definition introduces an abbreviation.
- ▶ Whenever we say “set”, we could instead say “an unordered collection of distinct objects” and vice versa.
- ▶ Definitions can also introduce notation.

German: Definition

# Disproofs

- ▶ A **disproof** (**refutation**) shows that a given mathematical statement is **false** by giving an example where the preconditions are true, but the conclusion is false.
- ▶ This requires deriving, in a sequence of proof steps, the opposite (negation) of the conclusion.

## Example (False statement)

“If  $p \in \mathbb{N}_0$  is a prime number then  $p$  is odd.”

## Refutation.

Consider natural number 2 as a counter example. It is prime because it has exactly 2 divisors, 1 and itself. It is not odd, because it is divisible by 2. □

German: Widerlegung

## Exercise

You want to disprove the following statement with a counterexample:

If the sun is shining then all kids eat ice cream.

What properties must your counterexample have?

[Discuss with your neighbour; 2 minutes]



# A Word on Style

A proof should help the reader to see why the result must be true.

- ▶ A proof should be easy to follow.
- ▶ Omit unnecessary information.
- ▶ Move self-contained parts into separate lemmas.
- ▶ In complicated proofs, reveal the overall structure in advance.
- ▶ Have a clear line of argument.

→ Writing a proof is like writing an essay.

# Proof Strategies

typical proof/disproof strategies:

- 1 “All  $x \in S$  with the property  $P$  also have the property  $Q$ .”  
“For all  $x \in S$ : if  $x$  has property  $P$ , then  $x$  has property  $Q$ .”
  - ▶ To prove, assume you are given an arbitrary  $x \in S$  that has the property  $P$ .  
Give a sequence of proof steps showing that  $x$  must have the property  $Q$ .
  - ▶ To disprove, find a **counterexample**, i. e., find an  $x \in S$  that has property  $P$  but not  $Q$  and prove this.

# Proof Strategies

typical proof/disproof strategies:

- ② “ $A$  is a subset of  $B$ .”
  - ▶ To prove, assume you have an arbitrary element  $x \in A$  and prove that  $x \in B$ .
  - ▶ To disprove, find an element in  $x \in A \setminus B$  and prove that  $x \in A \setminus B$ .

# Proof Strategies

typical proof/disproof strategies:

- ③ “For all  $x \in S$ :  $x$  has property  $P$  iff  $x$  has property  $Q$ .”  
(“iff”: “if and only if”)
  - ▶ To prove, separately prove “if  $P$  then  $Q$ ” and “if  $Q$  then  $P$ ”.
  - ▶ To disprove, disprove “if  $P$  then  $Q$ ” or disprove “if  $Q$  then  $P$ ”.

# Proof Strategies

typical proof/disproof strategies:

- ④ “ $A = B$ ”, where  $A$  and  $B$  are sets.
  - ▶ To prove, separately prove “ $A \subseteq B$ ” and “ $B \subseteq A$ ”.
  - ▶ To disprove, disprove “ $A \subseteq B$ ” or disprove “ $B \subseteq A$ ”.



# Proof Techniques

proof techniques we use in this course:

- ▶ direct proof
- ▶ indirect proof (proof by contradiction)
- ▶ structural induction

## A3.2 Direct Proof

# Direct Proof

## Direct Proof

Direct derivation of the statement by deducing or rewriting.

German: Direkter Beweis

## Direct Proof: Example

### Theorem (distributivity)

For all sets  $A, B, C$ :  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

### Proof.

We first show that  $x \in A \cap (B \cup C)$  implies  $x \in (A \cap B) \cup (A \cap C)$  ( $\subseteq$  part):

Let  $x \in A \cap (B \cup C)$ . Then by the definition of  $\cap$  it holds that  $x \in A$  and  $x \in B \cup C$ .

We make a case distinction between  $x \in B$  and  $x \notin B$ :

If  $x \in B$  then, because  $x \in A$  is true,  $x \in A \cap B$  must be true.

Otherwise, because  $x \in B \cup C$  we know that  $x \in C$  and thus with  $x \in A$ , that  $x \in A \cap C$ .

In both cases  $x \in A \cap B$  or  $x \in A \cap C$ ,  
and we conclude  $x \in (A \cap B) \cup (A \cap C)$ .

...

## Direct Proof: Example

### Theorem (distributivity)

For all sets  $A, B, C$ :  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

### Proof (continued).

$\supseteq$  part: we must show that  $x \in (A \cap B) \cup (A \cap C)$  implies  $x \in A \cap (B \cup C)$ .

Let  $x \in (A \cap B) \cup (A \cap C)$ .

We make a case distinction between  $x \in A \cap B$  and  $x \notin A \cap B$ :

If  $x \in A \cap B$  then  $x \in A$  and  $x \in B$ .

The latter implies  $x \in B \cup C$  and hence  $x \in A \cap (B \cup C)$ .

If  $x \notin A \cap B$  we know  $x \in A \cap C$  due to  $x \in (A \cap B) \cup (A \cap C)$ .

This (analogously) implies  $x \in A$  and  $x \in C$ , and hence  $x \in B \cup C$  and thus  $x \in A \cap (B \cup C)$ .

In both cases we conclude  $x \in A \cap (B \cup C)$ .

...

## Direct Proof: Example

### Theorem (distributivity)

For all sets  $A, B, C$ :  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

### Proof (continued).

We have shown that every element of  $A \cap (B \cup C)$  is an element of  $(A \cap B) \cup (A \cap C)$  and vice versa.

Thus, both sets are equal. □

## Direct Proof: Example

### Theorem (distributivity)

For all sets  $A, B, C$ :  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

### Proof.

Alternative:

$$\begin{aligned} A \cap (B \cup C) &= \{x \mid x \in A \text{ and } x \in B \cup C\} \\ &= \{x \mid x \in A \text{ and } (x \in B \text{ or } x \in C)\} \\ &= \{x \mid (x \in A \text{ and } x \in B) \text{ or } (x \in A \text{ and } x \in C)\} \\ &= \{x \mid x \in A \cap B \text{ or } x \in A \cap C\} \\ &= (A \cap B) \cup (A \cap C) \end{aligned}$$



## A3.3 Indirect Proof



# Indirect Proof

## Indirect Proof (Proof by Contradiction)

- ▶ Make an **assumption** that the statement is false.
- ▶ Use the assumption to derive a **contradiction**.
- ▶ This shows that the assumption must be false and hence the original statement must be true.

German: Indirekter Beweis, Beweis durch Widerspruch

## Indirect Proof: Example

### Theorem

Let  $A$  and  $B$  be sets. If  $A \setminus B = \emptyset$  then  $A \subseteq B$ .

### Proof.

We prove the theorem by contradiction.

Assume that there are sets  $A$  and  $B$  with  $A \setminus B = \emptyset$  and  $A \not\subseteq B$ .

Let  $A$  and  $B$  be such sets.

Since  $A \not\subseteq B$  there is some  $x \in A$  such that  $x \notin B$ .

For this  $x$  it holds that  $x \in A \setminus B$ .

This is a contradiction to  $A \setminus B = \emptyset$ .

We conclude that the assumption was false and thus the theorem is true. □

## A3.4 Structural Induction

## Inductively Defined Sets: Examples

### Example (Natural Numbers)

The set  $\mathbb{N}_0$  of natural numbers is inductively defined as follows:

- ▶ 0 is a natural number.
- ▶ If  $n$  is a natural number, then  $n + 1$  is a natural number.

### Example (Binary Tree)

The set  $\mathcal{B}$  of binary trees is inductively defined as follows:

- ▶  $\square$  is a binary tree (a leaf)
- ▶ If  $L$  and  $R$  are binary trees, then  $\langle L, \bigcirc, R \rangle$  is a binary tree (with inner node  $\bigcirc$ ).

**Implicit statement:** all elements of the set can be constructed by finite application of these rules

# Inductive Definition of a Set

## Inductive Definition

A set  $M$  can be defined **inductively** by specifying

- ▶ **basic elements** that are contained in  $M$
- ▶ **construction rules** of the form  
“Given some elements of  $M$ , another element of  $M$  can be constructed like this.”

German: Induktive Definition, Basiselemente, Konstruktionsregeln

# Structural Induction

## Structural Induction

Proof of statement for all elements of an inductively defined set

- ▶ **basis**: proof of the statement for the basic elements
- ▶ **induction hypothesis (IH)**:  
suppose that the statement is true for some elements  $M$
- ▶ **inductive step**: proof of the statement for elements constructed by applying a construction rule to  $M$   
(one inductive step for each construction rule)

German: Strukturelle Induktion, Induktionsanfang,  
Induktionsvoraussetzung, Induktionsschritt

## Structural Induction: Example (1)

### Definition (Leaves of a Binary Tree)

The number of **leaves** of a binary tree  $B$ , written  $leaves(B)$ , is defined as follows:

$$leaves(\square) = 1$$

$$leaves(\langle L, \circ, R \rangle) = leaves(L) + leaves(R)$$

### Definition (Inner Nodes of a Binary Tree)

The number of **inner nodes** of a binary tree  $B$ , written  $inner(B)$ , is defined as follows:

$$inner(\square) = 0$$

$$inner(\langle L, \circ, R \rangle) = inner(L) + inner(R) + 1$$

## Structural Induction: Example (2)

### Theorem

*For all binary trees  $B$ :  $inner(B) = leaves(B) - 1$ .*

### Proof.

induction basis:

$$inner(\square) = 0 = 1 - 1 = leaves(\square) - 1$$

$\rightsquigarrow$  statement is true for base case

...



## Structural Induction: Example (3)

Proof (continued).

induction hypothesis:

to prove that the statement is true for a composite tree  $\langle L, \circlearrowleft, R \rangle$ , we may use that it is true for the subtrees  $L$  and  $R$ .

inductive step for  $B = \langle L, \circlearrowleft, R \rangle$ :

$$\begin{aligned} \mathit{inner}(B) &= \mathit{inner}(L) + \mathit{inner}(R) + 1 \\ &\stackrel{\text{IH}}{=} (\mathit{leaves}(L) - 1) + (\mathit{leaves}(R) - 1) + 1 \\ &= \mathit{leaves}(L) + \mathit{leaves}(R) - 1 = \mathit{leaves}(B) - 1 \end{aligned}$$

□

## Structural Induction: Exercise (if time)

### Definition (Height of a Binary Tree)

The **height** of a binary tree  $B$ , written  $height(B)$ , is defined as follows:

$$height(\square) = 0$$

$$height(\langle L, \circlearrowleft, R \rangle) = \max\{height(L), height(R)\} + 1$$

Prove by structural induction:

### Theorem

For all binary trees  $B$ :  $leaves(B) \leq 2^{height(B)}$ .



## A3.5 Summary

# Summary

- ▶ A **proof** is based on axioms and previously proven statements.
- ▶ Individual **proof steps** must be obvious derivations.
- ▶ **direct proof**: sequence of derivations or rewriting
- ▶ **indirect proof**: refute the negated statement
- ▶ **structural induction**: generalization of mathematical induction to arbitrary recursive structures