

# Algorithms and Data Structures

## A6. Runtime Analysis: Logarithm


Gabriele Röger and Patrick Schneider

University of Basel

March 5, 2026

# Logarithm

# Content of the Course




# Logarithm

- For the analysis of merge sort, we will need the **logarithm function**.
- This is often the case in runtime analysis, in particular for divide-and-conquer algorithms.
- The logarithm to the base  $b$  is the inverse function to exponentiation with base  $b$ , i.e.

$$\log_b x = y \text{ iff. } b^y = x.$$

- **Example:**  $\log_2 8 = 3$ , because  $2^3 = 8$   
**Example:**  $\log_3 81 = 4$ , because  $3^4 = 81$
- $\log_b a$  intuitively (if this works without remainder):  
“How often must we divide  $a$  by  $b$  to reach 1?”

# Logarithm: Illustration



# Calculation with Logarithms

The following rules are immediate results of the rules  
 $(b^c)^d = b^{cd} = (b^c)^d$  and  $b^c b^d = b^{c+d}$ :

# Calculation with Logarithms

The following rules are immediate results of the rules  $(b^c)^d = b^{cd} = (b^c)^d$  and  $b^c b^d = b^{c+d}$ :

product  $\log_b(xy) = \log_b x + \log_b y$

# Calculation with Logarithms

The following rules are immediate results of the rules  $(b^c)^d = b^{cd} = (b^c)^d$  and  $b^c b^d = b^{c+d}$ :

product	$\log_b(xy) = \log_b x + \log_b y$
power	$\log_b(x^r) = r \log_b x$

# Calculation with Logarithms

The following rules are immediate results of the rules  $(b^c)^d = b^{cd} = (b^c)^d$  and  $b^c b^d = b^{c+d}$ :

product	$\log_b(xy) = \log_b x + \log_b y$
power	$\log_b(x^r) = r \log_b x$
change of base	$\log_b x = \log_a x / \log_a b$

## Logarithm: Example Calculation

In the analysis of algorithms, we sometimes see expressions of the form  $a^{\log_b x}$ . How do we get the logarithm out of the exponent?

## Logarithm: Example Calculation

In the analysis of algorithms, we sometimes see expressions of the form  $a^{\log_b x}$ . How do we get the logarithm out of the exponent?

Example:  $5^{\log_2 x}$

We use  $5 = 2^{\log_2 5}$ .

## Logarithm: Example Calculation

In the analysis of algorithms, we sometimes see expressions of the form  $a^{\log_b x}$ . How do we get the logarithm out of the exponent?

Example:  $5^{\log_2 x}$

We use  $5 = 2^{\log_2 5}$ .

$$\begin{aligned}5^{\log_2 x} &= (2^{\log_2 5})^{\log_2 x} \\ &= 2^{\log_2 5 \log_2 x} \\ &= 2^{\log_2 x \log_2 5} \\ &= (2^{\log_2 x})^{\log_2 5} \\ &= x^{\log_2 5} \\ &\approx x^{2.32}\end{aligned}$$