

# Discrete Mathematics in Computer Science

## B8. Functions

Malte Helmert, Gabriele Röger

University of Basel

October 19, 2020

# Discrete Mathematics in Computer Science

October 19, 2020 — B8. Functions

## B8.1 Partial and Total Functions

## B8.2 Operations on Partial Functions

## B8.3 Properties of Functions

## B8.1 Partial and Total Functions

## Important Building Blocks of Discrete Mathematics

Important building blocks:

- ▶ sets
- ▶ relations
- ▶ functions



In principle, functions are just a special kind of relations:

- ▶  $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $f(x) = x^2$
- ▶ relation  $R$  over  $\mathbb{N}_0$  with  $R = \{(x, y) \mid x, y \in \mathbb{N}_0 \text{ and } y = x^2\}$ .

## Functional Relations

### Definition

A binary relation  $R$  over sets  $A$  and  $B$  is **functional** if for every  $a \in A$  there is at most one  $b \in B$  with  $(a, b) \in R$ .



## Functions – Examples

►  $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $f(x) = x^2 + 1$

►  $abs : \mathbb{Z} \rightarrow \mathbb{N}_0$  with

$$abs(x) = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{otherwise} \end{cases}$$

►  $distance : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$  with

$$distance((x_1, y_1), (x_2, y_2)) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

## Partial Function – Example

Partial function  $r : \mathbb{Z} \times \mathbb{Z} \not\rightarrow \mathbb{Q}$  with

$$r(n, d) = \begin{cases} \frac{n}{d} & \text{if } d \neq 0 \\ \text{undefined} & \text{otherwise} \end{cases}$$

## Partial Functions

### Definition (Partial function)

A **partial function**  $f$  from set  $A$  to set  $B$  (written  $f : A \not\rightarrow B$ ) is given by a **functional relation**  $G$  over  $A$  and  $B$ .

Relation  $G$  is called the **graph** of  $f$ .

We write  $f(x) = y$  for  $(x, y) \in G$  and say  $y$  is the **image** of  $x$  under  $f$ .

If there is no  $y \in B$  with  $(x, y) \in G$ , then  $f(x)$  is **undefined**.

Partial function  $r : \mathbb{Z} \times \mathbb{Z} \not\rightarrow \mathbb{Q}$  with

$$r(n, d) = \begin{cases} \frac{n}{d} & \text{if } d \neq 0 \\ \text{undefined} & \text{otherwise} \end{cases}$$

has graph  $\{( (n, d), \frac{n}{d} ) \mid n \in \mathbb{Z}, d \in \mathbb{Z} \setminus \{0\} \} \subseteq \mathbb{Z}^2 \times \mathbb{Q}$ .

## Domain (of Definition), Codomain, Image

### Definition (domain of definition, codomain, image)

Let  $f : A \rightarrow B$  be a partial function.


Set  $A$  is called the **domain** of  $f$ , set  $B$  is its **codomain**.

The **domain of definition** of  $f$  is the set

$\text{dom}(f) = \{x \in A \mid \text{there is a } y \in B \text{ with } f(x) = y\}$ .

The **image (or range)** of  $f$  is the set

$\text{img}(f) = \{y \mid \text{there is an } x \in A \text{ with } f(x) = y\}$ .



$f : \{a, b, c, d, e\} \rightarrow \{1, 2, 3, 4\}$   
 $f(a) = 4, f(b) = 2, f(c) = 1, f(e) = 4$   
 domain  $\{a, b, c, d, e\}$   
 codomain  $\{1, 2, 3, 4\}$   
 domain of definition  $\text{dom}(f) = \{a, b, c, e\}$   
 image  $\text{img}(f) = \{1, 2, 4\}$

## Preimage


The preimage contains all elements of the domain that are mapped to given elements of the codomain.

### Definition (Preimage)

Let  $f : A \rightarrow B$  be a partial function and let  $Y \subseteq B$ .

The **preimage of  $Y$  under  $f$**  is the set

$f^{-1}[Y] = \{x \in A \mid f(x) \in Y\}$ .




$f^{-1}[\{1\}] =$   
 $f^{-1}[\{3\}] =$   
 $f^{-1}[\{4\}] =$   
 $f^{-1}[\{1, 2\}] =$

## Total Functions

### Definition (Total function)

A **(total) function**  $f : A \rightarrow B$  from set  $A$  to set  $B$  is a partial function from  $A$  to  $B$  such that  $f(x)$  is defined for all  $x \in A$ .

→ no difference between the domain and the domain of definition



## Specifying a Function

Some common ways of specifying a function:

- ▶ Listing the mapping **explicitly**, e. g.  
 $f(a) = 4, f(b) = 2, f(c) = 1, f(e) = 4$  or  
 $f = \{a \mapsto 4, b \mapsto 2, c \mapsto 1, e \mapsto 4\}$
- ▶ By a **formula**, e. g.  $f(x) = x^2 + 1$
- ▶ By **recurrence**, e. g.  
 $0! = 1$  and  
 $n! = n(n-1)!$  for  $n > 0$
- ▶ In terms of other functions, e. g. inverse, composition

## Relationship to Functions in Programming

```
def factorial(n):
    if n == 0:
        return 1
    else:
        return n * factorial(n-1)
```

→ Relationship between recursion and recurrence

## Relationship to Functions in Programming

```
def foo(n):
    value = ...
    while <some condition>:
        ...
        value = ...
    return value
```

→ Does possibly not terminate on all inputs.  
 → Value is undefined for such inputs.  
 → Theoretical computer science: partial function

## Relationship to Functions in Programming

```
import random
counter = 0

def bar(n):
    print("Hi! I got input", n)
    global counter
    counter += 1
    return random.choice([1,2,n])
```

→ Functions in programming don't always compute mathematical functions (except *purely functional languages*).  
 → In addition, not all mathematical functions are computable.

## B8.2 Operations on Partial Functions

## Restrictions and Extensions

### Definition (restriction and extension)

Let  $f : A \rightarrow B$  be a partial function and let  $X \subseteq A$ .

The **restriction of  $f$  to  $X$**  is the partial function  $f|_X : X \rightarrow B$  with  $f|_X(x) = f(x)$  for all  $x \in X$ .

A function  $f' : A' \rightarrow B$  is called an **extension of  $f$**  if  $A \subseteq A'$  and  $f'|_A = f$ .

The restriction of  $f$  to its domain of definition is a total function.

What's the graph of the restriction?

What's the restriction of  $f$  to its domain?

## Function Composition

### Definition (Composition of partial functions)

Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be partial functions.

The **composition of  $f$  and  $g$**  is  $g \circ f : A \rightarrow C$  with

$$(g \circ f)(x) = \begin{cases} g(f(x)) & \text{if } f \text{ is defined for } x \text{ and} \\ & g \text{ is defined for } f(x) \\ \text{undefined} & \text{otherwise} \end{cases}$$

Corresponds to relation composition of the graphs.

If  $f$  and  $g$  are functions, their composition is a function.

Example:

$$f : \mathbb{N}_0 \rightarrow \mathbb{N}_0 \quad \text{with } f(x) = x^2$$

$$g : \mathbb{N}_0 \rightarrow \mathbb{N}_0 \quad \text{with } g(x) = x + 3$$

$$(g \circ f)(x) =$$

## Properties of Function Composition

Function composition is

- ▶ **not commutative:**
  - ▶  $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $f(x) = x^2$
  - ▶  $g : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $g(x) = x + 3$
  - ▶  $(g \circ f)(x) = x^2 + 3$
  - ▶  $(f \circ g)(x) = (x + 3)^2$
- ▶ **associative**, i.e.  $h \circ (g \circ f) = (h \circ g) \circ f$   
→ analogous to associativity of relation composition

## Function Composition in Programming

We implicitly compose functions all the time...

```
def foo(n):
    ...
    x = somefunction(n)
    y = someotherfunction(x)
    ...

```

Many languages also allow explicit composition of functions,  
e.g. in Haskell:

```
incr x = x + 1
square x = x * x
squareplusone = incr . square
```

## B8.3 Properties of Functions

## Properties of Functions


- ▶ Partial functions map every element of their domain to at most one element of their codomain, total functions map it to exactly one such value.
- ▶ Different elements of the domain can have the same image.
- ▶ There can be values of the codomain that aren't the image of any element of the domain.
- ▶ We often want to exclude such cases  
→ define additional properties to say this quickly

### Injective Functions


An **injective function** maps distinct elements of its domain to distinct elements of its co-domain.

#### Definition (Injective Function)

A function  $f : A \rightarrow B$  is **injective** (also **one-to-one** or an **injection**) if for all  $x, y \in A$  with  $x \neq y$  it holds that  $f(x) \neq f(y)$ .



injective



not injective

### Injective Functions – Examples

Which of these functions are injective?

- ▶  $f : \mathbb{Z} \rightarrow \mathbb{N}_0$  with  $f(x) = |x|$
- ▶  $g : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $g(x) = x^2$
- ▶  $h : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $h(x) = \begin{cases} x - 1 & \text{if } x \text{ is odd} \\ x + 1 & \text{if } x \text{ is even} \end{cases}$

## Composition of Injective Functions

### Theorem

If  $f : A \rightarrow B$  and  $g : B \rightarrow C$  are injective functions then also  $g \circ f$  is injective.

### Proof.

Consider arbitrary elements  $x, y \in A$  with  $x \neq y$ .

Since  $f$  is injective, we know that  $f(x) \neq f(y)$ .

As  $g$  is injective, this implies that  $g(f(x)) \neq g(f(y))$ .

With the definition of  $g \circ f$ , we conclude that

$(g \circ f)(x) \neq (g \circ f)(y)$ .


Overall, this shows that  $g \circ f$  is injective.  $\square$

## Surjective Functions


A **surjective function** maps at least one elements to every element of its co-domain.

### Definition (Surjective Function)

A function  $f : A \rightarrow B$  is **surjective** (also **onto** or a **surjection**) if its **image is equal to its codomain**, i.e. for all  $y \in B$  there is an  $x \in A$  with  $f(x) = y$ .



surjective



not surjective

## Surjective Functions – Examples

Which of these functions are surjective?

- ▶  $f : \mathbb{Z} \rightarrow \mathbb{N}_0$  with  $f(x) = |x|$
- ▶  $g : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $g(x) = x^2$
- ▶  $h : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $h(x) = \begin{cases} x - 1 & \text{if } x \text{ is odd} \\ x + 1 & \text{if } x \text{ is even} \end{cases}$

## Composition of Surjective Functions

### Theorem

If  $f : A \rightarrow B$  and  $g : B \rightarrow C$  are surjective functions then also  $g \circ f$  is surjective.

### Proof.

Consider an arbitrary element  $z \in C$ .

Since  $g$  is surjective, there is a  $y \in B$  with  $g(y) = z$ .

As  $f$  is surjective, for such a  $y$  there is an  $x \in A$  with  $f(x) = y$  and thus  $g(f(x)) = z$ .


Overall, for every  $z \in C$  there is an  $x \in A$  with  $(g \circ f)(x) = g(f(x)) = z$ , so  $g \circ f$  is surjective.  $\square$

## Bijective Functions

A **bijective function** pairs every element of its domain with exactly one element of its codomain and every element of the codomain is paired with exactly one element of the domain.

### Definition (Bijective Function)

A function is **bijective** (also a **one-to-one correspondence** or a **bijection**) if it is **injective** and **surjective**.



### Corollary

*The composition of two bijective functions is bijective.*

## Bijective Functions – Examples

Which of these functions are bijective?



- ▶  $f : \mathbb{Z} \rightarrow \mathbb{N}_0$  with  $f(x) = |x|$
- ▶  $g : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $g(x) = x^2$
- ▶  $h : \mathbb{N}_0 \rightarrow \mathbb{N}_0$  with  $h(x) = \begin{cases} x - 1 & \text{if } x \text{ is odd} \\ x + 1 & \text{if } x \text{ is even} \end{cases}$

## Inverse Function

### Definition

Let  $f : A \rightarrow B$  be a bijection.

The **inverse function** of  $f$  is the function  $f^{-1} : B \rightarrow A$  with  $f^{-1}(y) = x$  iff  $f(x) = y$ .



## Inverse Function and Composition

### Theorem

Let  $f : A \rightarrow B$  be a bijection.

- ① For all  $x \in A$  it holds that  $f^{-1}(f(x)) = x$ .
- ② For all  $y \in B$  it holds that  $f(f^{-1}(y)) = y$ .
- ③  $(f^{-1})^{-1} = f$

### Proof sketch.

- ① For  $x \in A$  let  $y = f(x)$ . Then  $f^{-1}(f(x)) = f^{-1}(y) = x$
- ② For  $y \in B$  there is exactly one  $x$  with  $y = f(x)$ . With this  $x$  it holds that  $f^{-1}(y) = x$  and overall  $f(f^{-1}(y)) = f(x) = y$ .
- ③ Def. of inverse:  $(f^{-1})^{-1}(x) = y$  iff  $f^{-1}(y) = x$  iff  $f(x) = y$ .

## Inverse Function

### Theorem

Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be bijections.

Then  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ .

### Proof.

We need to show that for all  $x \in C$  it holds that

$$(g \circ f)^{-1}(x) = (f^{-1} \circ g^{-1})(x).$$

Consider an arbitrary  $x \in C$  and let  $y = (g \circ f)^{-1}(x)$ .

By the definition of the inverse  $(g \circ f)(y) = x$ .

Let  $z = f(y)$ . With  $(g \circ f)(y) = g(f(y))$ , we know that  $x = g(z)$ .

From  $z = f(y)$  we get  $f^{-1}(z) = y$  and

from  $x = g(z)$  we get  $g^{-1}(x) = z$ .

This gives  $(f^{-1} \circ g^{-1})(x) = f^{-1}(g^{-1}(x)) = f^{-1}(z) = y$ . □

## Summary

- ▶ **injective function**: maps distinct elements of its domain to distinct elements of its co-domain.
- ▶ **surjective function**: maps at least one elements to every element of its co-domain.
- ▶ **bijective function**: injective and surjective  
→ one-to-one correspondence
- ▶ Bijective functions are invertible. The **inverse** function of  $f$  maps the image of  $x$  under  $f$  to  $x$ .