

# Discrete Mathematics in Computer Science

## Equivalence Relations and Partitions

Malte Helmert, Gabriele Röger

University of Basel

## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .

## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .
- Possible properties of homogeneous relations  $R$  over  $S$ :

## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .
- Possible properties of homogeneous relations  $R$  over  $S$ :
  - **reflexive**:  $(x, x) \in R$  for all  $x \in S$

## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .
- Possible properties of homogeneous relations  $R$  over  $S$ :
  - **reflexive**:  $(x, x) \in R$  for all  $x \in S$
  - **irreflexive**:  $(x, x) \notin R$  for all  $x \in S$

## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .
- Possible properties of homogeneous relations  $R$  over  $S$ :
  - **reflexive**:  $(x, x) \in R$  for all  $x \in S$
  - **irreflexive**:  $(x, x) \notin R$  for all  $x \in S$
  - **symmetric**:  $(x, y) \in R$  iff  $(y, x) \in R$

## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .
- Possible properties of homogeneous relations  $R$  over  $S$ :
  - **reflexive**:  $(x, x) \in R$  for all  $x \in S$
  - **irreflexive**:  $(x, x) \notin R$  for all  $x \in S$
  - **symmetric**:  $(x, y) \in R$  iff  $(y, x) \in R$
  - **asymmetric**: if  $(x, y) \in R$  then  $(y, x) \notin R$

## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .
- Possible properties of homogeneous relations  $R$  over  $S$ :
  - **reflexive**:  $(x, x) \in R$  for all  $x \in S$
  - **irreflexive**:  $(x, x) \notin R$  for all  $x \in S$
  - **symmetric**:  $(x, y) \in R$  iff  $(y, x) \in R$
  - **asymmetric**: if  $(x, y) \in R$  then  $(y, x) \notin R$
  - **antisymmetric**: if  $(x, y) \in R$  then  $(y, x) \notin R$  or  $x = y$



## Relations: Recap

- A **relation over sets**  $S_1, \dots, S_n$  is a set  $R \subseteq S_1 \times \dots \times S_n$ .
- Possible properties of homogeneous relations  $R$  over  $S$ :
  - **reflexive**:  $(x, x) \in R$  for all  $x \in S$
  - **irreflexive**:  $(x, x) \notin R$  for all  $x \in S$
  - **symmetric**:  $(x, y) \in R$  iff  $(y, x) \in R$
  - **asymmetric**: if  $(x, y) \in R$  then  $(y, x) \notin R$
  - **antisymmetric**: if  $(x, y) \in R$  then  $(y, x) \notin R$  or  $x = y$
  - **transitive**: if  $(x, y) \in R$  and  $(y, z) \in R$  then  $(x, z) \in R$

# Motivation

- Think of any attribute that two objects can have in common, e. g. their color.
- We could place the objects into distinct “buckets”, e. g. one bucket for each color.
- We also can define a relation  $\sim$  such that  $x \sim y$  iff  $x$  and  $y$  share the attribute, e. g. have the same color.
- Would this relation be
  - reflexive?
  - irreflexive?
  - symmetric?
  - asymmetric?
  - antisymmetric?
  - transitive?

# Equivalence Relation

## Definition (Equivalence Relation)

A binary relation  $\sim$  over set  $S$  is an **equivalence relation** if  $\sim$  is **reflexive, symmetric and transitive**.

Is this definition indeed what we want?

Does it allow us to partition the objects into buckets

(e. g. one group for all objects that share a specific color)?

# Partition

## Definition (Partition)

A **partition** of a set  $S$  is a set  $P \subseteq \mathcal{P}(S)$  such that

- $X \neq \emptyset$  for all  $X \in P$ ,
- $\bigcup_{X \in P} X = S$ , and
- $X \cap Y = \emptyset$  for all  $X, Y \in P$  with  $X \neq Y$ ,

The elements of  $P$  are called the **blocks** of the partition.

# Partition

Let  $S = \{e_1, \dots, e_5\}$ .

Which of the following sets are partitions of  $S$ ?

- $P_1 = \{\{e_1, e_4\}, \{e_3\}, \{e_2, e_5\}\}$

# Partition

Let  $S = \{e_1, \dots, e_5\}$ .

Which of the following sets are partitions of  $S$ ?

- $P_1 = \{\{e_1, e_4\}, \{e_3\}, \{e_2, e_5\}\}$
- $P_2 = \{\{e_1, e_4, e_5\}, \{e_3\}\}$

# Partition

Let  $S = \{e_1, \dots, e_5\}$ .

Which of the following sets are partitions of  $S$ ?

- $P_1 = \{\{e_1, e_4\}, \{e_3\}, \{e_2, e_5\}\}$
- $P_2 = \{\{e_1, e_4, e_5\}, \{e_3\}\}$
- $P_3 = \{\{e_1, e_4, e_5\}, \{e_3\}, \{e_2, e_5\}\}$

# Partition

Let  $S = \{e_1, \dots, e_5\}$ .

Which of the following sets are partitions of  $S$ ?

- $P_1 = \{\{e_1, e_4\}, \{e_3\}, \{e_2, e_5\}\}$
- $P_2 = \{\{e_1, e_4, e_5\}, \{e_3\}\}$
- $P_3 = \{\{e_1, e_4, e_5\}, \{e_3\}, \{e_2, e_5\}\}$
- $P_4 = \{\{e_1\}, \{e_2\}, \{e_3\}, \{e_4\}, \{e_5\}\}$



# Partition

Let  $S = \{e_1, \dots, e_5\}$ .

Which of the following sets are partitions of  $S$ ?

- $P_1 = \{\{e_1, e_4\}, \{e_3\}, \{e_2, e_5\}\}$
- $P_2 = \{\{e_1, e_4, e_5\}, \{e_3\}\}$
- $P_3 = \{\{e_1, e_4, e_5\}, \{e_3\}, \{e_2, e_5\}\}$
- $P_4 = \{\{e_1\}, \{e_2\}, \{e_3\}, \{e_4\}, \{e_5\}\}$
- $P_5 = \{\{e_1\}, \{e_2\}, \{e_3\}, \{e_4\}, \{e_5\}, \{\}\}$

# A Property of Partitions

## Lemma

*Let  $S$  be a set and  $P$  be a partition of  $S$ .*

*Then every  $x \in S$  is an element of exactly one  $X \in P$ .*

**Proof:**  $\rightsquigarrow$  exercises

## Block of an Element

The lemma enables the following definition:

### Definition

Let  $S$  be a set and  $P$  be a partition of  $S$ .

For  $e \in S$  we denote by  $[e]_P$  the block  $X \in P$  such that  $e \in X$ .

## Block of an Element

The lemma enables the following definition:

### Definition

Let  $S$  be a set and  $P$  be a partition of  $S$ .

For  $e \in S$  we denote by  $[e]_P$  the block  $X \in P$  such that  $e \in X$ .

Consider partition  $P = \{\{e_1, e_4\}, \{e_3\}, \{e_2, e_5\}\}$  of  $\{e_1, \dots, e_5\}$ .

$[e_1]_P =$

# Connection between Partitions and Equivalence Relations?

- We will now explore the connection between partitions and equivalence relations.
- **Spoiler:** They are essentially the same concept.

# Partitions Induce Equivalence Relations I

## Definition (Relation induced by a partition)

Let  $S$  be a set and  $P$  be a partition of  $S$ .

The relation  $\sim_P$  induced by  $P$  is the binary relation over  $S$  with

$$x \sim_P y \text{ iff } [x]_P = [y]_P.$$

$x \sim_P y$  iff  $x$  and  $y$  are in the same block of  $P$ .

# Partitions Induce Equivalence Relations I

## Definition (Relation induced by a partition)

Let  $S$  be a set and  $P$  be a partition of  $S$ .

The relation  $\sim_P$  induced by  $P$  is the binary relation over  $S$  with

$$x \sim_P y \text{ iff } [x]_P = [y]_P.$$

$x \sim_P y$  iff  $x$  and  $y$  are in the same block of  $P$ .

Consider partition  $P = \{\{1, 4, 5\}, \{2, 3\}\}$  of set  $\{1, 2, \dots, 5\}$ .

$$\sim_P = \{(1, 1), (1, 4), (1, 5), (4, 1), (4, 4), (4, 5), (5, 1), (5, 4), (5, 5), \\ (2, 2), (2, 3), (3, 2), (3, 3)\}$$

We will show that  $\sim_P$  is an equivalence relation.

## Partitions Induce Equivalence Relations II

### Theorem

Let  $P$  be a partition of  $S$ .

Relation  $\sim_P$  induced by  $P$  is an *equivalence relation* over  $S$ .



## Partitions Induce Equivalence Relations II

### Theorem

Let  $P$  be a partition of  $S$ .

Relation  $\sim_P$  induced by  $P$  is an *equivalence relation* over  $S$ .

### Proof.

We need to show that  $\sim_P$  is reflexive, symmetric and transitive.

## Partitions Induce Equivalence Relations II

### Theorem

Let  $P$  be a partition of  $S$ .

Relation  $\sim_P$  induced by  $P$  is an *equivalence relation* over  $S$ .

### Proof.

We need to show that  $\sim_P$  is reflexive, symmetric and transitive.

**reflexive:** As  $=$  is reflexive it holds for all  $x \in S$  that  $[x]_P = [x]_P$  and hence also that  $x \sim_P x$ .

## Partitions Induce Equivalence Relations II

### Theorem

Let  $P$  be a partition of  $S$ .

Relation  $\sim_P$  induced by  $P$  is an **equivalence relation** over  $S$ .

### Proof.

We need to show that  $\sim_P$  is reflexive, symmetric and transitive.

**reflexive:** As  $=$  is reflexive it holds for all  $x \in S$  that  $[x]_P = [x]_P$  and hence also that  $x \sim_P x$ .

**symmetric:** If  $x \sim_P y$  then  $[x]_P = [y]_P$ . With the symmetry of  $=$  we get that  $[y]_P = [x]_P$  and conclude that  $y \sim_P x$ .

## Partitions Induce Equivalence Relations II

### Theorem

Let  $P$  be a partition of  $S$ .

Relation  $\sim_P$  induced by  $P$  is an **equivalence relation** over  $S$ .

### Proof.

We need to show that  $\sim_P$  is reflexive, symmetric and transitive.

**reflexive:** As  $=$  is reflexive it holds for all  $x \in S$  that  $[x]_P = [x]_P$  and hence also that  $x \sim_P x$ .

**symmetric:** If  $x \sim_P y$  then  $[x]_P = [y]_P$ . With the symmetry of  $=$  we get that  $[y]_P = [x]_P$  and conclude that  $y \sim_P x$ .

**transitive:** If  $x \sim_P y$  and  $y \sim_P z$  then  $[x]_P = [y]_P$  and  $[y]_P = [z]_P$ . As  $=$  is transitive, it then also holds that  $[x]_P = [z]_P$  and hence  $x \sim_P z$ . □

# Equivalence Classes

## Definition (equivalence class)

Let  $R$  be an equivalence relation over set  $S$ .

For any  $x \in S$ , the **equivalence class of  $x$**  is the set

$$[x]_R = \{y \in S \mid xRy\}.$$

# Equivalence Classes

## Definition (equivalence class)

Let  $R$  be an equivalence relation over set  $S$ .

For any  $x \in S$ , the **equivalence class of  $x$**  is the set

$$[x]_R = \{y \in S \mid xRy\}.$$

Consider

$$R = \{(1, 1), (1, 4), (1, 5), (4, 1), (4, 4), (4, 5), (5, 1), (5, 4), (5, 5), \\ (2, 2), (2, 3), (3, 2), (3, 3)\}$$

over set  $\{1, 2, \dots, 5\}$ .

$$[4]_R =$$

# Equivalence Relations Induce Partitions

## Theorem

Let  $R$  be an equivalence relation over set  $S$ .

The set  $P = \{[x]_R \mid x \in S\}$  is a *partition of  $S$* .

- 1) For  $x \in S$ , it holds that  $x \in [x]_R$  because  $R$  is reflexive.  
Hence, no  $X \in P$  is empty.

...

# Equivalence Relations Induce Partitions

## Theorem

Let  $R$  be an equivalence relation over set  $S$ .

The set  $P = \{[x]_R \mid x \in S\}$  is a *partition of  $S$* .

## Proof.

We need to show that

- 1  $X \neq \emptyset$  for all  $X \in P$ ,
- 2  $\bigcup_{X \in P} X = S$ , and
- 3  $X \cap Y = \emptyset$  for all  $X, Y \in P$  with  $X \neq Y$ ,



# Equivalence Relations Induce Partitions

## Theorem

Let  $R$  be an equivalence relation over set  $S$ .

The set  $P = \{[x]_R \mid x \in S\}$  is a *partition of  $S$* .

## Proof.

We need to show that

- 1)  $X \neq \emptyset$  for all  $X \in P$ ,
- 2)  $\bigcup_{X \in P} X = S$ , and
- 3)  $X \cap Y = \emptyset$  for all  $X, Y \in P$  with  $X \neq Y$ ,

1) For  $x \in S$ , it holds that  $x \in [x]_R$  because  $R$  is reflexive.

Hence, no  $X \in P$  is empty. ...

## Equivalence Relations Induce Partitions

Proof (continued).

For 2) we show  $\bigcup_{X \in P} X \subseteq S$  and  $\bigcup_{X \in P} X \supseteq S$  separately.

## Equivalence Relations Induce Partitions

Proof (continued).

For 2) we show  $\bigcup_{X \in P} X \subseteq S$  and  $\bigcup_{X \in P} X \supseteq S$  separately.

$\subseteq$ : Consider an arbitrary  $x \in \bigcup_{X \in P} X$ . Since  $x$  is contained in the union, it must be an element of some  $X \in P$ . Consider such an  $X$ . By the definition of  $P$ , there is a  $y \in S$  such that  $X = [y]_R$ .

Since  $x \in [y]_R$ , it holds that  $yRx$ .

As  $R$  is a relation over  $S$ , this implies that  $x \in S$ .

## Equivalence Relations Induce Partitions

Proof (continued).

For 2) we show  $\bigcup_{X \in P} X \subseteq S$  and  $\bigcup_{X \in P} X \supseteq S$  separately.

$\subseteq$ : Consider an arbitrary  $x \in \bigcup_{X \in P} X$ . Since  $x$  is contained in the union, it must be an element of some  $X \in P$ . Consider such an  $X$ . By the definition of  $P$ , there is a  $y \in S$  such that  $X = [y]_R$ .

Since  $x \in [y]_R$ , it holds that  $yRx$ .

As  $R$  is a relation over  $S$ , this implies that  $x \in S$ .

$\supseteq$ : Consider an arbitrary  $x \in S$ . Since  $x \in [x]_R$  (cf. 1) and  $[x]_R \in P$ , it holds that  $x \in \bigcup_{X \in P} X$ .

...

# Equivalence Relations Induce Partitions

Proof (continued).

We show 3) by contrapositive:

For all  $X, Y \in P$ : if  $X \cap Y \neq \emptyset$  then  $X = Y$ .

## Equivalence Relations Induce Partitions

Proof (continued).

We show 3) by contrapositive:

For all  $X, Y \in P$ : if  $X \cap Y \neq \emptyset$  then  $X = Y$ .

Let  $X, Y$  be two sets from  $P$  with  $X \cap Y \neq \emptyset$ .

Then there is an  $e$  with  $e \in X \cap Y$  and there are  $x, y \in S$  with  $X = [x]_R$  and  $Y = [y]_R$ . Consider such  $e, x, y$ .

## Equivalence Relations Induce Partitions

Proof (continued).

We show 3) by contrapositive:

For all  $X, Y \in P$ : if  $X \cap Y \neq \emptyset$  then  $X = Y$ .

Let  $X, Y$  be two sets from  $P$  with  $X \cap Y \neq \emptyset$ .

Then there is an  $e$  with  $e \in X \cap Y$  and there are  $x, y \in S$  with  $X = [x]_R$  and  $Y = [y]_R$ . Consider such  $e, x, y$ .

As  $e \in [x]_R$  and  $e \in [y]_R$  it holds that  $xRe$  and  $yRe$ . Since  $R$  is symmetric, we get from  $yRe$  that  $eRy$ . By transitivity,  $xRe$  and  $eRy$  imply  $xRy$ , which by symmetry also gives  $yRx$ .

## Equivalence Relations Induce Partitions

Proof (continued).

We show 3) by contrapositive:

For all  $X, Y \in P$ : if  $X \cap Y \neq \emptyset$  then  $X = Y$ .

Let  $X, Y$  be two sets from  $P$  with  $X \cap Y \neq \emptyset$ .

Then there is an  $e$  with  $e \in X \cap Y$  and there are  $x, y \in S$  with  $X = [x]_R$  and  $Y = [y]_R$ . Consider such  $e, x, y$ .

As  $e \in [x]_R$  and  $e \in [y]_R$  it holds that  $xRe$  and  $yRe$ . Since  $R$  is symmetric, we get from  $yRe$  that  $eRy$ . By transitivity,  $xRe$  and  $eRy$  imply  $xRy$ , which by symmetry also gives  $yRx$ .

We show  $[x]_R \subseteq [y]_R$ : consider an arbitrary  $z \in [x]_R$ . Then  $xRz$ . From  $yRx$  and  $xRz$ , by transitivity we get  $yRz$ . This establishes  $z \in [y]_R$ . As  $z$  was chosen arbitrarily, it holds that  $[x]_R \subseteq [y]_R$ .



## Equivalence Relations Induce Partitions

Proof (continued).

We show 3) by contrapositive:

For all  $X, Y \in P$ : if  $X \cap Y \neq \emptyset$  then  $X = Y$ .

Let  $X, Y$  be two sets from  $P$  with  $X \cap Y \neq \emptyset$ .

Then there is an  $e$  with  $e \in X \cap Y$  and there are  $x, y \in S$  with  $X = [x]_R$  and  $Y = [y]_R$ . Consider such  $e, x, y$ .

As  $e \in [x]_R$  and  $e \in [y]_R$  it holds that  $xRe$  and  $yRe$ . Since  $R$  is symmetric, we get from  $yRe$  that  $eRy$ . By transitivity,  $xRe$  and  $eRy$  imply  $xRy$ , which by symmetry also gives  $yRx$ .

We show  $[x]_R \subseteq [y]_R$ : consider an arbitrary  $z \in [x]_R$ . Then  $xRz$ . From  $yRx$  and  $xRz$ , by transitivity we get  $yRz$ . This establishes  $z \in [y]_R$ . As  $z$  was chosen arbitrarily, it holds that  $[x]_R \subseteq [y]_R$ .

Analogously, we can show that  $[x]_R \supseteq [y]_R$ , so overall  $X = Y$ .  $\square$

# Summary

- We typically encounter equivalence relations when we consider objects as equivalent wrt. some attribute/property.
- A relation is an **equivalence relation** if it is **reflexive, symmetric and transitive**.
- A **partition** of a set groups the elements into non-empty subsets.
- The concepts are closely connected:  
in principle just different perspectives on the same “situation”.

# Discrete Mathematics in Computer Science

## Partial Orders

Malte Helmert, Gabriele Röger

University of Basel

# Order Relations

- An **equivalence relation** is **reflexive, symmetric and transitive**.
- Such a relation induces a partition into “equivalent” objects.

# Order Relations

- An **equivalence relation** is **reflexive, symmetric and transitive**.
- Such a relation induces a partition into “equivalent” objects.
- We now consider **other combinations of properties**, that allow us to **compare objects** in a set against other objects.

# Order Relations

- An **equivalence relation** is **reflexive, symmetric and transitive**.
- Such a relation induces a partition into “equivalent” objects.
- We now consider **other combinations of properties**, that allow us to **compare objects** in a set against other objects.
- “Number  $x$  is not larger than number  $y$ .”  
“Set  $S$  is a subset of set  $T$ .”  
“Jerry runs at least as fast as Tom.”  
“Pasta tastes better than Potatoes.”

# Partial Orders

- We begin with **partial orders**.

# Partial Orders

- We begin with **partial orders**.
- Example partial order relations are  $\leq$  over  $\mathbb{N}$  or  $\subseteq$  for sets.



# Partial Orders

- We begin with **partial orders**.
- Example partial order relations are  $\leq$  over  $\mathbb{N}$  or  $\subseteq$  for sets.
- Are these relations
  - reflexive?
  - irreflexive?
  - symmetric?
  - asymmetric?
  - antisymmetric?
  - transitive?

## Partial Orders – Definition

Definition (Partial order, partially ordered sets)

A binary relation  $\preceq$  over set  $S$  is a **partial order** if  $\preceq$  is **reflexive, antisymmetric and transitive**.

## Partial Orders – Definition

### Definition (Partial order, partially ordered sets)

A binary relation  $\preceq$  over set  $S$  is a **partial order** if  $\preceq$  is **reflexive, antisymmetric and transitive**.

A **partially ordered set** (or **poset**) is a pair  $(S, R)$  where  $S$  is a **set** and  $R$  is a **partial order over  $S$** .

## Partial Orders – Definition

### Definition (Partial order, partially ordered sets)

A binary relation  $\preceq$  over set  $S$  is a **partial order** if  $\preceq$  is **reflexive, antisymmetric and transitive**.

A **partially ordered set** (or **poset**) is a pair  $(S, R)$  where  $S$  is a **set** and  $R$  is a **partial order over  $S$** .

Which of these relations are partial orders?

- strict subset relation  $\subset$  for sets
- not-less-than relation  $\geq$  over  $\mathbb{N}_0$
- $R = \{(a, a), (a, b), (b, b), (b, c), (c, c)\}$  over  $\{a, b, c\}$

# Least and Greatest Element

Some special elements of posets:

## Definition (Least and greatest element)

Let  $\preceq$  be a partial order over set  $S$ .

An element  $x \in S$  is the **least element** of  $S$  if **for all**  $y \in S$  it holds that  $x \preceq y$ .

It is the **greatest element** of  $S$  if **for all**  $y \in S$ ,  $y \preceq x$ .

# Least and Greatest Element

Some special elements of posets:

## Definition (Least and greatest element)

Let  $\preceq$  be a partial order over set  $S$ .

An element  $x \in S$  is the **least element** of  $S$  if **for all**  $y \in S$  it holds that  $x \preceq y$ .

It is the **greatest element** of  $S$  if **for all**  $y \in S$ ,  $y \preceq x$ .

- Is there a least/greatest element? Which one?
  - $S = \{1, 2, 3\}$  and  $\preceq = \{(x, y) \mid x, y \in S \text{ and } x \leq y\}$ .

# Least and Greatest Element

Some special elements of posets:

## Definition (Least and greatest element)

Let  $\preceq$  be a partial order over set  $S$ .

An element  $x \in S$  is the **least element** of  $S$  if **for all**  $y \in S$  it holds that  $x \preceq y$ .

It is the **greatest element** of  $S$  if **for all**  $y \in S$ ,  $y \preceq x$ .

- Is there a least/greatest element? Which one?
  - $S = \{1, 2, 3\}$  and  $\preceq = \{(x, y) \mid x, y \in S \text{ and } x \leq y\}$ .
  - $\mathbb{N}_0$  and standard relation  $\leq$ .

# Least and Greatest Element

Some special elements of posets:

## Definition (Least and greatest element)

Let  $\preceq$  be a partial order over set  $S$ .

An element  $x \in S$  is the **least element** of  $S$  if **for all**  $y \in S$  it holds that  $x \preceq y$ .

It is the **greatest element** of  $S$  if **for all**  $y \in S$ ,  $y \preceq x$ .

- Is there a least/greatest element? Which one?
  - $S = \{1, 2, 3\}$  and  $\preceq = \{(x, y) \mid x, y \in S \text{ and } x \leq y\}$ .
  - $\mathbb{N}_0$  and standard relation  $\leq$ .
- Why can we say **the** least element instead of **a** least element?



# Uniqueness of Least Element

## Theorem

*Let  $\preceq$  be a partial order over set  $S$ .*

*If  $S$  contains a least element, it contains exactly one least element.*

# Uniqueness of Least Element

## Theorem

Let  $\preceq$  be a partial order over set  $S$ .

If  $S$  contains a least element, it contains exactly one least element.

## Proof.

By contradiction: Assume  $x, y$  are least elements of  $S$  with  $x \neq y$ .



# Uniqueness of Least Element

## Theorem

Let  $\preceq$  be a partial order over set  $S$ .

If  $S$  contains a least element, it contains exactly one least element.

## Proof.

**By contradiction:** Assume  $x, y$  are least elements of  $S$  with  $x \neq y$ .

Since  $x$  is a least element,  $x \preceq y$  is true.

Since  $y$  is a least element,  $y \preceq x$  is true.



# Uniqueness of Least Element

## Theorem

Let  $\preceq$  be a partial order over set  $S$ .

If  $S$  contains a least element, it contains exactly one least element.

## Proof.

**By contradiction:** Assume  $x, y$  are least elements of  $S$  with  $x \neq y$ .

Since  $x$  is a least element,  $x \preceq y$  is true.

Since  $y$  is a least element,  $y \preceq x$  is true.

As a partial order is antisymmetric, this implies that  $x = y$ .  $\downarrow$  □

# Uniqueness of Least Element

## Theorem

Let  $\preceq$  be a partial order over set  $S$ .

If  $S$  contains a least element, it contains exactly one least element.

## Proof.

**By contradiction:** Assume  $x, y$  are least elements of  $S$  with  $x \neq y$ .

Since  $x$  is a least element,  $x \preceq y$  is true.

Since  $y$  is a least element,  $y \preceq x$  is true.

As a partial order is antisymmetric, this implies that  $x = y$ .  $\downarrow$   $\square$

Analogously: If there is a greatest element then is unique.

## Minimal and Maximal Elements

### Definition (Minimal/Maximal element of a set)

Let  $\preceq$  be a partial order over set  $S$ .

An element  $x \in S$  is a **minimal element** of  $S$  if **there is no  $y \in S$  with  $y \preceq x$  and  $x \neq y$ .**

An element  $x \in S$  is a **maximal element** of  $S$  if **there is no  $y \in S$  with  $x \preceq y$  and  $x \neq y$ .**

## Minimal and Maximal Elements

### Definition (Minimal/Maximal element of a set)

Let  $\preceq$  be a partial order over set  $S$ .

An element  $x \in S$  is a **minimal element** of  $S$  if **there is no  $y \in S$  with  $y \preceq x$  and  $x \neq y$ .**

An element  $x \in S$  is a **maximal element** of  $S$  if **there is no  $y \in S$  with  $x \preceq y$  and  $x \neq y$ .**

A set can have several minimal elements and no least element.

Example?

# Total Orders

- Relations  $\leq$  over  $\mathbb{N}_0$  and  $\subseteq$  for sets are partial orders.



# Total Orders

- Relations  $\leq$  over  $\mathbb{N}_0$  and  $\subseteq$  for sets are partial orders.
- Can we compare every object against every object?

# Total Orders

- Relations  $\leq$  over  $\mathbb{N}_0$  and  $\subseteq$  for sets are partial orders.
- Can we compare every object against every object?
  - For all  $x, y \in \mathbb{N}_0$  it holds that  $x \leq y$  or that  $y \leq x$  (or both).

# Total Orders

- Relations  $\leq$  over  $\mathbb{N}_0$  and  $\subseteq$  for sets are partial orders.
- Can we compare every object against every object?
  - For all  $x, y \in \mathbb{N}_0$  it holds that  $x \leq y$  or that  $y \leq x$  (or both).
  - $\{1, 2\} \not\subseteq \{2, 3\}$  and  $\{2, 3\} \not\subseteq \{1, 2\}$

# Total Orders

- Relations  $\leq$  over  $\mathbb{N}_0$  and  $\subseteq$  for sets are partial orders.
- Can we compare every object against every object?
  - For all  $x, y \in \mathbb{N}_0$  it holds that  $x \leq y$  or that  $y \leq x$  (or both).
  - $\{1, 2\} \not\subseteq \{2, 3\}$  and  $\{2, 3\} \not\subseteq \{1, 2\}$
- Relation  $\leq$  is a **total** order, relation  $\subseteq$  is not.

## Total Order – Definition

### Definition (Total relation)

A binary relation  $R$  over set  $S$  is **total** (or **connex**) if for all  $x, y \in S$  at least one of  $xRy$  or  $yRx$  is true.

## Total Order – Definition

### Definition (Total relation)

A binary relation  $R$  over set  $S$  is **total** (or **connex**) if for all  $x, y \in S$  at least one of  $xRy$  or  $yRx$  is true.

### Definition (Total order)

A binary relation is a **total order** if it is **total** and a **partial order**.

# Summary

- A **partial order** is **reflexive, antisymmetric and transitive**.
- With a **total order**  $\preceq$  over  $S$  there are no elements  $x, y \in S$  with  $x \not\preceq y$  and  $y \not\preceq x$ .
- If  $x$  is the **greatest element** of a set  $S$ , it is greater than every element: for all  $y \in S$  it holds that  $y \preceq x$ .
- If  $x$  is a **maximal element** of set  $S$  then it is not smaller than any other element  $y$ : there is no  $y \in S$  with  $x \preceq y$  and  $x \neq y$ .

# Discrete Mathematics in Computer Science

## Strict Orders

Malte Helmert, Gabriele Röger

University of Basel



# Strict Orders

- A **partial** order is reflexive, antisymmetric and transitive.
- We now consider **strict orders**.

# Strict Orders

- A **partial** order is reflexive, antisymmetric and transitive.
- We now consider **strict orders**.
- Example strict order relations are  $<$  over  $\mathbb{N}$  or  $\subset$  for sets.

# Strict Orders

- A **partial** order is reflexive, antisymmetric and transitive.
- We now consider **strict orders**.
- Example strict order relations are  $<$  over  $\mathbb{N}$  or  $\subset$  for sets.
- Are these relations
  - reflexive?
  - irreflexive?
  - symmetric?
  - asymmetric?
  - antisymmetric?
  - transitive?

## Strict Orders – Definition

### Definition (Strict order)

A binary relation  $\prec$  over set  $S$  is a **strict order** if  $\prec$  is **irreflexive, asymmetric and transitive**.

## Strict Orders – Definition

### Definition (Strict order)

A binary relation  $\prec$  over set  $S$  is a **strict order** if  $\prec$  is **irreflexive, asymmetric and transitive**.

Which of these relations are strict orders?

- subset relation  $\subseteq$  for sets
- strict superset relation  $\supset$  for sets

## Strict Orders – Definition

### Definition (Strict order)

A binary relation  $\prec$  over set  $S$  is a **strict order** if  $\prec$  is **irreflexive, asymmetric and transitive**.

Which of these relations are strict orders?

- subset relation  $\subseteq$  for sets
- strict superset relation  $\supset$  for sets

Can a relation be both, a partial order and a strict order?

## Strict Total Orders

- As partial orders, a strict order does not automatically allow us to rank arbitrary two objects against each other.

# Strict Total Orders

- As partial orders, a strict order does not automatically allow us to rank arbitrary two objects against each other.

- **Example 1** (personal preferences):

- “Pasta tastes better than potato.”


- “Rice tastes better than bread.”

- “Bread tastes better than potato.”

- “Rice tastes better than potato.”

- This definition of “tastes better than” is a strict order.

- No ranking of pasta against rice or of pasta against bread.






# Strict Total Orders

- As partial orders, a strict order does not automatically allow us to rank arbitrary two objects against each other.

- **Example 1** (personal preferences):

- “Pasta tastes better than potato.”
- “Rice tastes better than bread.”
- “Bread tastes better than potato.”
- “Rice tastes better than potato.”
- This definition of “tastes better than” is a strict order.
- No ranking of pasta against rice or of pasta against bread.




- **Example 2:**  $\subset$  relation for sets

# Strict Total Orders

- As partial orders, a strict order does not automatically allow us to rank arbitrary two objects against each other.

- **Example 1** (personal preferences):

- “Pasta tastes better than potato.”
- “Rice tastes better than bread.”
- “Bread tastes better than potato.”
- “Rice tastes better than potato.”
- This definition of “tastes better than” is a strict order.
- No ranking of pasta against rice or of pasta against bread.



- **Example 2:**  $\subset$  relation for sets

- It **doesn't work** to simply require that the strict order is total.  
Why?

## Strict Total Orders – Definition

### Definition (Trichotomy)

A binary relation  $R$  over set  $S$  is **trichotomous** if for all  $x, y \in S$  exactly one of  $xRy$ ,  $yRx$  or  $x = y$  is true.

## Strict Total Orders – Definition

### Definition (Trichotomy)

A binary relation  $R$  over set  $S$  is **trichotomous** if for all  $x, y \in S$  exactly one of  $xRy$ ,  $yRx$  or  $x = y$  is true.

### Definition (Strict total order)

A binary relation  $\prec$  over  $S$  is a **strict total order** if  $\prec$  is **trichotomous** and a **strict order**.

## Strict Total Orders – Definition

### Definition (Trichotomy)

A binary relation  $R$  over set  $S$  is **trichotomous** if for all  $x, y \in S$  exactly one of  $xRy$ ,  $yRx$  or  $x = y$  is true.

### Definition (Strict total order)

A binary relation  $\prec$  over  $S$  is a **strict total order** if  $\prec$  is **trichotomous** and a **strict order**.

A strict total order completely ranks the elements of set  $S$ .

**Example:**  $<$  relation over  $\mathbb{N}_0$  gives the standard ordering  $0, 1, 2, 3, \dots$  of natural numbers.

# Special Elements

Special elements are defined almost as for partial orders:

Definition (Least/greatest/minimal/maximal element of a set)

Let  $\prec$  be a **strict** order over set  $S$ .

An element  $x \in S$  is the **least element** of  $S$   
if for all  $y \in S$  where  $y \neq x$  it holds that  $x \prec y$ .


It is the **greatest element** of  $S$  if for all  $y \in S$  where  $y \neq x$ ,  $y \prec x$ .

Element  $x \in S$  is a **minimal element** of  $S$   
if there is no  $y \in S$  with  $y \prec x$ .

It is a **maximal element** of  $S$   
if there is no  $y \in S$  with  $x \prec y$ .

## Special Elements – Example

Consider again the previous example:

$$S = \{\text{Pasta, Potato, Bread, Rice}\}$$
$$\prec = \{(\text{Pasta, Potato}), (\text{Bread, Potato}), \\ (\text{Rice, Potato}), (\text{Rice, Bread})\}$$


Is there a least and a greatest element?

Which elements are maximal or minimal?

# Summary and Outlook

- A **strict order** is **irreflexive**, **asymmetric** and **transitive**.
- Strict **total** orders and **special elements** are analogously defined as for partial sets but with a special treatment of equal elements.
- For partial order  $\preceq$  we can define a related strict order  $\prec$  as  $x \prec y$  if  $x \preceq y$  and  $y \not\preceq x$ .
- For strict order  $\prec$  we can define a related partial order  $\preceq$  as  $x \preceq y$  if  $x \prec y$  or  $x = y$ .
- There are more related concepts, e. g.
  - **(total) preorder**: (connex), reflexive, transitive
  - **well-order**: total order over  $S$  such that every non-empty subset has a least element