

Discrete Mathematics in Computer Science

Cardinality of Infinite Sets

Malte Helmert, Gabriele Röger

University of Basel

Finite Sets Revisited

We already know:

- The **cardinality** $|S|$ measures the size of set S .
- A set is **finite** if it has a finite number of elements.
- The **cardinality** of a finite set is the **number of elements** it contains.
- For a finite set S , it holds that $|\mathcal{P}(S)| = 2^{|S|}$.

Finite Sets Revisited

We already know:

- The **cardinality** $|S|$ measures the size of set S .
- A set is **finite** if it has a finite number of elements.
- The **cardinality** of a finite set is the **number of elements** it contains.
- For a finite set S , it holds that $|\mathcal{P}(S)| = 2^{|S|}$.

A set is **infinite** if it has an infinite number of elements.

- Do all infinite sets have the same cardinality?
- Does the power set of infinite set S have the same cardinality as S ?

Comparing the Cardinality of Sets

- $\{1, 2, 3\}$ and $\{\text{dog, cat, mouse}\}$ have cardinality 3.
- We can pair their elements:

1 \leftrightarrow dog

2 \leftrightarrow cat

3 \leftrightarrow mouse

Comparing the Cardinality of Sets

- $\{1, 2, 3\}$ and $\{\text{dog, cat, mouse}\}$ have cardinality 3.
- We can pair their elements:

1 \leftrightarrow dog

2 \leftrightarrow cat

3 \leftrightarrow mouse

- We call such a mapping a **bijection** from one set to the other.
 - Each element of one set is paired with exactly one element of the other set.
 - Each element of the other set is paired with exactly one element of the first set.

Equinumerous Sets

We use the existence of a pairing also as criterion for infinite sets:

Definition (Equinumerous Sets)

Two sets A and B have the same cardinality ($|A| = |B|$) if there **exists a bijection from A to B** .

Such sets are called **equinumerous**.

Equinumerous Sets

We use the existence of a pairing also as criterion for infinite sets:

Definition (Equinumerous Sets)

Two sets A and B have the same cardinality ($|A| = |B|$) if there **exists a bijection from A to B** .

Such sets are called **equinumerous**.

When is a set “smaller” than another set?

Comparing the Cardinality of Sets

- Consider $A = \{1, 2\}$ and $B = \{\text{dog}, \text{cat}, \text{mouse}\}$.
- We can map distinct elements of A to distinct elements of B :

$1 \mapsto \text{dog}$

$2 \mapsto \text{cat}$

- We call this an **injective function** from A to B :
 - every element of A is mapped to an element of B ;
 - different elements of A are mapped to different elements of B .

Comparing Cardinality

Definition (cardinality not larger)

Set A has **cardinality less than or equal** to the cardinality of set B ($|A| \leq |B|$), if **there is an injective function from A to B** .

Definition (strictly smaller cardinality)

Set A has **cardinality strictly less** than the cardinality of set B ($|A| < |B|$), if $|A| \leq |B|$ and $|A| \neq |B|$.

Comparing Cardinality

Definition (cardinality not larger)

Set A has **cardinality less than or equal** to the cardinality of set B ($|A| \leq |B|$), if **there is an injective function from A to B** .

Definition (strictly smaller cardinality)

Set A has **cardinality strictly less** than the cardinality of set B ($|A| < |B|$), if $|A| \leq |B|$ and $|A| \neq |B|$.

Consider set A and object $e \notin A$. Is $|A| < |A \cup \{e\}|$?

Discrete Mathematics in Computer Science

Hilbert's Hotel

Malte Helmert, Gabriele Röger

University of Basel


Hilbert's Hotel

Our intuition for finite sets does not always work for infinite sets.

Hilbert's Hotel

Our intuition for finite sets does not always work for infinite sets.


- If in a hotel all rooms are occupied then it cannot accommodate additional guests.



Hilbert's Hotel

Our intuition for finite sets does not always work for infinite sets.

- If in a hotel all rooms are occupied then it cannot accommodate additional guests.
- But **Hilbert's Grand Hotel** has **infinitely many rooms**.
- All these rooms are **occupied**.




One More Guest Arrives




- Every guest moves from her current room n to room $n + 1$.
- Room 1 is then free.
- The new guest gets room 1.

Four More Guests Arrive



- Every guest moves from her current room n to room $n + 4$.
- Rooms 1 to 4 are no longer occupied and can be used for the new guests.


Four More Guests Arrive




- Every guest moves from her current room n to room $n + 4$.
- Rooms 1 to 4 are no longer occupied and can be used for the new guests.

→ Works for any finite number of additional guests.

An Infinite Number of Guests Arrives



An Infinite Number of Guests Arrives



- Every guest moves from her current room n to room $2n$.
- The infinitely many rooms with odd numbers are now available.
- The new guests fit into these rooms.

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests

... arrive?

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests
- infinitely many ferries, each with an infinite number of coaches, each with infinitely many guests

... arrive?

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests
- infinitely many ferries, each with an infinite number of coaches, each with infinitely many guests
- ...

... arrive?

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests
- infinitely many ferries, each with an infinite number of coaches, each with infinitely many guests
- ...

... arrive?

There are strategies for all these situations as long as with “infinite” we mean “countably infinite” and there is a finite number of layers.

Discrete Mathematics in Computer Science

\aleph_0 and Countable Sets

Malte Helmert, Gabriele Röger

University of Basel

Comparing Cardinality

- Two sets A and B have the **same cardinality** if their elements can be paired (i.e. there is a bijection from A to B).
- Set A has a **strictly smaller cardinality** than set B if
 - we can map distinct elements of A to distinct elements of B (i.e. there is an injective function from A to B), and
 - $|A| \neq |B|$.

Comparing Cardinality

- Two sets A and B have the **same cardinality** if their elements can be paired (i.e. there is a bijection from A to B).
- Set A has a **strictly smaller cardinality** than set B if
 - we can map distinct elements of A to distinct elements of B (i.e. there is an injective function from A to B), and
 - $|A| \neq |B|$.
- This clearly makes sense for finite sets.

Comparing Cardinality

- Two sets A and B have the **same cardinality** if their elements can be paired (i.e. there is a bijection from A to B).
- Set A has a **strictly smaller cardinality** than set B if
 - we can map distinct elements of A to distinct elements of B (i.e. there is an injective function from A to B), and
 - $|A| \neq |B|$.
- This clearly makes sense for finite sets.
- What about infinite sets?
Do they even have different cardinalities?

The Cardinality of the Natural Numbers

Definition (\aleph_0)

The **cardinality of \mathbb{N}_0** is denoted by \aleph_0 , i.e. $\aleph_0 = |\mathbb{N}_0|$

Read: “aleph-zero”, “aleph-nought” or “aleph-null”

Countable and Countably Infinite Sets

Definition (countably infinite and countable)

A set A is **countably infinite** if $|A| = |\mathbb{N}_0|$.

A set A is **countable** if $|A| \leq |\mathbb{N}_0|$.

A set is **countable** if it is **finite or countably infinite**.

Countable and Countably Infinite Sets

Definition (countably infinite and countable)

A set A is **countably infinite** if $|A| = |\mathbb{N}_0|$.

A set A is **countable** if $|A| \leq |\mathbb{N}_0|$.

A set is **countable** if it is **finite or countably infinite**.

- We can count the elements of a countable set one at a time.
- The objects are “**discrete**” (in contrast to “**continuous**”).
- **Discrete mathematics** deals with all kinds of countable sets.

Set of Even Numbers

- $even = \{n \mid n \in \mathbb{N}_0 \text{ and } n \text{ is even}\}$
- Obviously: $even \subset \mathbb{N}_0$
- Intuitively, there are twice as many natural numbers as even numbers — no?
- Is $|even| < |\mathbb{N}_0|$?

Set of Even Numbers

Theorem (set of even numbers is countably infinite)

The set of all even natural numbers is countably infinite, i. e. $|\{n \mid n \in \mathbb{N}_0 \text{ and } n \text{ is even}\}| = |\mathbb{N}_0|$.

Set of Even Numbers

Theorem (set of even numbers is countably infinite)

The set of all even natural numbers is countably infinite, i. e. $|\{n \mid n \in \mathbb{N}_0 \text{ and } n \text{ is even}\}| = |\mathbb{N}_0|$.

Proof Sketch.

We can pair every natural number n with the even number $2n$. \square

Set of Perfect Squares

Theorem (set of perfect squares is countably infinite)

*The set of all perfect squares is countably infinite,
i. e. $|\{n^2 \mid n \in \mathbb{N}_0\}| = |\mathbb{N}_0|$.*

Set of Perfect Squares

Theorem (set of perfect squares is countably infinite)

*The set of all perfect squares is countably infinite,
i. e. $|\{n^2 \mid n \in \mathbb{N}_0\}| = |\mathbb{N}_0|$.*

Proof Sketch.

We can pair every natural number n with square number n^2 . □

Subsets of Countable Sets are Countable

In general:

Theorem (subsets of countable sets are countable)

Let A be a countable set. Every set B with $B \subseteq A$ is countable.

Subsets of Countable Sets are Countable

In general:

Theorem (subsets of countable sets are countable)

Let A be a countable set. Every set B with $B \subseteq A$ is countable.

Proof.

Since A is countable there is an injective function f from A to \mathbb{N}_0 .
The restriction of f to B is an injective function from B to \mathbb{N}_0 . \square

Set of the Positive Rationals

Theorem (set of positive rationals is countably infinite)


Set $\mathbb{Q}_+ = \{n \mid n \in \mathbb{Q} \text{ and } n > 0\} = \{p/q \mid p, q \in \mathbb{N}_1\}$
is *countably infinite*.

Set of the Positive Rationals

Theorem (set of positive rationals is countably infinite)

Set $\mathbb{Q}_+ = \{n \mid n \in \mathbb{Q} \text{ and } n > 0\} = \{p/q \mid p, q \in \mathbb{N}_1\}$
is *countably infinite*.

Proof idea.



Union of Two Countable Sets is Countable

Theorem (union of two countable sets countable)

Let A and B be countable sets. Then $A \cup B$ is countable.

Proof sketch.

As A and B are countable there is an injective function f_A from A to \mathbb{N}_0 , analogously f_B from B to \mathbb{N}_0 .

We define function $f_{A \cup B}$ from $A \cup B$ to \mathbb{N}_0 as

$$f_{A \cup B}(e) = \begin{cases} 2f_A(e) & \text{if } e \in A \\ 2f_B(e) + 1 & \text{otherwise} \end{cases}$$

This $f_{A \cup B}$ is an injective function from $A \cup B$ to \mathbb{N}_0 . □

Integers and Rationals

Theorem (sets of integers and rationals are countably infinite)

The sets \mathbb{Z} and \mathbb{Q} are *countably infinite*.

Without proof (\rightsquigarrow exercises)

Union of More than Two Sets

Definition (arbitrary unions)

Let M be a set of sets. The union $\bigcup_{S \in M} S$ is the set with

$$x \in \bigcup_{S \in M} S \text{ iff exists } S \in M \text{ with } x \in S.$$

Countable Union of Countable Sets

Theorem

Let M be a *countable set of countable sets*.

Then $\bigcup_{S \in M}$ *is countable*.

We prove this formally after we have studied functions.

Set of all Binary Trees is Countable

Theorem (set of all binary trees is countable)

The set $B = \{b \mid b \text{ is a binary tree}\}$ is countable.

Proof.

For $n \in \mathbb{N}_0$ the set B_n of all binary trees with n leaves is finite.

With $M = \{B_i \mid i \in \mathbb{N}_0\}$ the set of all binary trees is

$$B = \bigcup_{B' \in M} B'.$$

Since M is a countable set of countable sets, B is countable. \square

And Now?

We have seen several sets with cardinality \aleph_0 .

And Now?

We have seen several sets with cardinality \aleph_0 .

What about our original questions?

- Do all infinite sets have the same cardinality?
- Does the power set of infinite set S have the same cardinality as S ?