Discrete Mathematics in Computer Science B4. Tuples & Cartesian Product

Malte Helmert, Gabriele Röger

University of Basel

Malte Helmert, Gabriele Röger (University of Discrete Mathematics in Computer Science

Malte Helmert, Gabriele Röger (University of Discrete Mathematics in Computer Science

Malte Helmert, Gabriele Röger (University of Discrete Mathematics in Computer Science

B4. Tuples & Cartesian Product

Tuples and the Cartesian Product

B4.1 Tuples and the Cartesian Product '

B4. Tuples & Cartesian Product

Tuples and the Cartesian Product

Sets vs. Tuples

- A set is an unordered collection of distinct objects.
- ► A tuple is an ordered sequence of objects.

Discrete Mathematics in Computer Science

B4.1 Tuples and the Cartesian Product

— B4. Tuples & Cartesian Product

3 / 8

Malte Helmert, Gabriele Röger (University of Discrete Mathematics in Computer Science

Tuples

- ▶ k-tuple: ordered sequence of k objects ($k \in \mathbb{N}_0$)
- ightharpoonup written (o_1, \ldots, o_k) or $\langle o_1, \ldots, o_k \rangle$
- ▶ unlike sets, order matters $(\langle 1, 2 \rangle \neq \langle 2, 1 \rangle)$
- objects may occur multiple times in a tuple
- objects contained in tuples are called components
- terminology:
 - k = 2: (ordered) pair
 - k = 3: triple
 - more rarely: quadruple, quintuple, sextuple, septuple, ...
- ▶ if k is clear from context (or does not matter), often just called tuple

Malte Helmert, Gabriele Röger (University of Discrete Mathematics in Computer Science

5 /

B4. Tuples & Cartesian Product

Tuples and the Cartesian Product

Cartesian Product

Definition (Cartesian Product and Cartesian Power)

Let S_1, \ldots, S_n be sets. The Cartesian product $S_1 \times \cdots \times S_n$ is the following set of *n*-tuples:

$$S_1 \times \cdots \times S_n = \{\langle x_1, \dots, x_n \rangle \mid x_1 \in S_1, x_2 \in S_2, \dots, x_n \in S_n \}.$$

The *k*-ary Cartesian power of a set *S* (with $k \in \mathbb{N}_1$) is the set $S^k = \{\langle o_1, \dots, o_k \rangle \mid o_i \in S \text{ for all } i \in \{1, \dots, k\}\} = \underbrace{S \times \dots \times S}_{k \text{ times}}.$

René Descartes: French mathematician and philosopher (1596-1650)

Example:
$$A = \{a, b\}, B = \{1, 2, 3\}$$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

 $A^2 = \{(a, a), (a, b), (b, a), (b, b)\}$

Equality of Tuples

B4. Tuples & Cartesian Product

Definition (Equality of Tuples)

Two *n*-tuples $t = \langle o_1, \ldots, o_n \rangle$ and $t' = \langle o'_1, \ldots, o'_n \rangle$ are equal (t = t') if for $i \in \{1, \ldots, n\}$ it holds that $o_i = o'_i$.

Malte Helmert, Gabriele Röger (University of Discrete Mathematics in Computer Science

6 /

B4. Tuples & Cartesian Product

Tuples and the Cartesian Product

(Non-)properties of the Cartesian Product

The Cartesian product is

- ▶ not commutative, in most cases $A \times B \neq B \times A$.
- ▶ not associative, in most cases $(A \times B) \times C \neq A \times (B \times C)$

Why? Exceptions?