## Discrete Mathematics in Computer Science C1. Introduction to Graphs

Malte Helmert, Gabriele Röger

University of Basel

# Discrete Mathematics in Computer Science — C1. Introduction to Graphs

#### C1.1 Graphs and Directed Graphs

## C1.2 Induced Graphs and Degree Lemma

# C1.1 Graphs and Directed Graphs

#### Graphs

Graphs (of various kinds) are ubiquitous in Computer Science and its applications.

Some examples:

- Boolean circuits in hardware design
- control flow graphs in compilers
- pathfinding in video games
- computer networks
- neural networks
- social networks

## Graph Theory

- Graph theory was founded in 1736 by Leonhard Euler's study of the Seven Bridges of Königsberg problem.
- It remains one of the main areas of discrete mathematics to this day.

More on Euler and the Seven Bridges of Königsberg:



#### The Seven Bridges of Königsberg – Numberphile. https://youtu.be/W18FDEA1jRQ

## Graphs and Directed Graphs – Definitions

Definition (Graph)
A graph (also: undirected graph) is a pair G = (V, E), where
V is a finite set called the set of vertices, and
E ⊆ {{u, v} ⊆ V | u ≠ v} is called the set of edges.

German: Graph, ungerichteter Graph, Knoten, Kanten

#### Definition (Directed Graph)

A directed graph (also: digraph) is a pair G = (N, A), where

N is a finite set called the set of nodes, and

• 
$$A \subseteq N \times N$$
 is called the set of arcs.

German: gerichteter Graph, Digraph, Knoten, Kanten/Pfeile

Graphs and Directed Graphs

#### Graphs and Directed Graphs – Pictorially

#### often described pictorially:



graph (V, E)



directed graph (N, A)

$$N = \{1, 2, 3, 4, 5\}$$

$$A = \{(1, 2), (1, 3), (2, 1), (3, 5), (4, 3), (4, 4), (5, 3), (5, 4)\}$$

#### Relationship to Relations

graphs vs. directed graphs:

- edges are sets of two elements, arcs are pairs
- arcs can be self-loops (v, v); edges cannot (why not?)

(di-)graphs vs. relations:

 A directed graph (N, A) is essentially identical to (= contains the same information as) an arbitrary relation R<sub>A</sub> over the finite set N: u R<sub>A</sub> v iff (u, v) ∈ A

A graph (V, E) is essentially identical to an irreflexive symmetric relation R<sub>E</sub> over the finite set V: u R<sub>E</sub> v iff {u, v} ∈ E

#### Other Kinds of Graphs

many variations exist, for example:

- self-loops may be allowed in edges ("non-simple" graphs)
- labeled graphs: additional information associated with vertices and/or edges
- weighted graphs: numbers associated with edges
- multigraphs: multiple edges between same vertices allowed
- mixed graphs: both edges and arcs allowed
- hypergraphs: edges can involve more than 2 vertices
- infinite graphs: may have infinitely many vertices/edges

# Graph Terminology

#### Definition (Graph Terminology)

Let (V, E) be a graph.

- *u* and *v* are the endpoints of the edge  $\{u, v\} \in E$
- *u* and *v* are incident to the edge  $\{u, v\} \in E$
- u and v are adjacent if  $\{u, v\} \in E$
- the vertices adjacent with v ∈ V are its neighbours neigh(v): neigh(v) = {w ∈ V | {v, w} ∈ E}
- ► the number of neighbours of v ∈ V is its degree deg(v): deg(v) = |neigh(v)|

# German: Endknoten, inzident, adjazent/benachbart, Nachbarn, Grad

#### Graph Terminology – Examples



endpoints, incident, adjacent, neighbours, degree

# Directed Graph Terminology

#### Definition (Directed Graph Terminology)

Let (N, A) be a directed graph.

- u is the tail and v is the head of the arc (u, v) ∈ A; we say (u, v) is an arc from u to v
- u and v are incident to the arc  $(u, v) \in A$
- *u* is a predecessor of *v* and *v* is a successor of *u* if  $(u, v) \in A$
- The predecessors and successor of v are written as pred(v) = {u ∈ N | (u, v) ∈ A} and succ(v) = {w ∈ N | (v, w) ∈ A}
- ► the number of predecessors/successors of v ∈ N is its indegree/outdegree: indeg(v) = |pred(v)|, outdeg(v) = |succ(v)|

# German: Fuss, Kopf, inzident, Vorgänger, Nachfolger, Eingangs-/Ausgangsgrad

Graphs and Directed Graphs

#### Directed Graph Terminology - Examples



head, tail, predecessors, successors, indegree, outdegree

# C1.2 Induced Graphs and Degree Lemma

## Induced Graph of a Directed Graph

Definition (undirected graph induced by a directed graph) Let G = (N, A) be a directed graph. The (undirected) graph induced by G is the graph (N, E) with  $E = \{\{u, v\} \mid (u, v) \in A, u \neq v\}.$ 

German: induziert

Questions:

- Why require  $u \neq v$ ?
- If |N| = n and |A| = m, how many vertices and edges does the induced graph have?
- ▶ How does the answer change if *G* has no self-loops?

Induced Graphs and Degree Lemma

#### Induced Graph of a Directed Graph – Example



#### Degree Lemma

Lemma (degree lemma for directed graphs) Let (N, A) be a directed graph. Then  $\sum_{v \in N} indeg(v) = \sum_{v \in N} outdeg(v) = |A|$ .

Intuitively: every arc contributes 1 to the indegree of one node and 1 to the outdegree of one node.

```
Lemma (degree lemma for undirected graphs)
Let (V, E) be a graph.
Then \sum_{v \in V} \deg(v) = 2|E|.
```

Intuitively: every edge contributes 1 to the degree of two vertices.

#### Corollary

Every graph has an even number of vertices with odd degree.

#### Degree Lemma – Example



$$\sum_{v \in V} \deg(v)$$
  
= deg(A) + deg(B) + deg(C) + deg(D) + deg(E) + deg(F) + deg(G)  
= 2 + 2 + 3 + 1 + 1 + 1 + 0

 $= 10 = 2 \cdot 5 = 2|E|$ 

#### 4 vertices with odd degree

## Degree Lemma – Proof (1)

Proof of degree lemma for directed graphs.

$$\sum_{v \in N} \operatorname{indeg}(v) = \sum_{v \in N} |\operatorname{pred}(v)|$$

$$= \sum_{v \in N} |\{u \mid u \in N, (u, v) \in A\}|$$

$$= \sum_{v \in N} |\{(u, v) \mid u \in N, (u, v) \in A\}|$$

$$= \left| \bigcup_{v \in N} \{(u, v) \mid u \in N, (u, v) \in A\} \right|$$

$$= |\{(u, v) \mid u \in N, v \in N, (u, v) \in A\}|$$

$$= |A|.$$

$$\sum_{v \in N} \operatorname{outdeg}(v) = |A| \text{ is analogous.}$$

# Degree Lemma – Proof (2)

We omit the proof for undirected graphs, which can be conducted similarly.

One possible proof strategy that reuses the result we proved:

- Define directed graph (V, A) from the graph (V, E) by orienting each edge into an arc arbitrarily.
- Observe deg(v) = indeg(v) + outdeg(v), where deg refers to the graph and indeg/outdeg to the directed graph.
- ► Use the degree lemma for directed graphs:  $\sum_{v \in V} \deg(v) = \sum_{v \in V} (\operatorname{indeg}(v) + \operatorname{outdeg}(v)) =$   $\sum_{v \in V} \operatorname{indeg}(v) + \sum_{v \in V} \operatorname{outdeg}(v) = |A| + |A| = 2|A| = 2|E|$