

Discrete Mathematics in Computer Science

Properties of Propositional Formulas

Malte Helmert, Gabriele Röger

University of Basel

The Story So Far

- **propositional logic** based on atomic propositions
- **syntax**: which formulas are well-formed?
- **semantics**: when is a formula true?
- **interpretations**: important basis of semantics

Reminder: Syntax of Propositional Logic

Definition (Syntax of Propositional Logic)

Let A be a set of **atomic propositions**. The set of **propositional formulas** (over A) is inductively defined as follows:

- Every **atom** $a \in A$ is a propositional formula over A .
- If φ is a propositional formula over A , then so is its **negation** $\neg\varphi$.
- If φ and ψ are propositional formulas over A , then so is the **conjunction** $(\varphi \wedge \psi)$.
- If φ and ψ are propositional formulas over A , then so is the **disjunction** $(\varphi \vee \psi)$.

The **implication** $(\varphi \rightarrow \psi)$ is an abbreviation for $(\neg\varphi \vee \psi)$.

The **biconditional** $(\varphi \leftrightarrow \psi)$ is an abbrev. for $((\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi))$.

Reminder: Semantics of Propositional Logic

Definition (Semantics of Propositional Logic)

A **truth assignment** (or **interpretation**) for a set of atomic propositions A is a function $\mathcal{I} : A \rightarrow \{0, 1\}$.

A propositional **formula** φ (over A) **holds under** \mathcal{I} (written as $\mathcal{I} \models \varphi$) according to the following definition:

$\mathcal{I} \models a$	iff	$\mathcal{I}(a) = 1$	(for $a \in A$)
$\mathcal{I} \models \neg\varphi$	iff	not $\mathcal{I} \models \varphi$	
$\mathcal{I} \models (\varphi \wedge \psi)$	iff	$\mathcal{I} \models \varphi$ and $\mathcal{I} \models \psi$	
$\mathcal{I} \models (\varphi \vee \psi)$	iff	$\mathcal{I} \models \varphi$ or $\mathcal{I} \models \psi$	

Properties of Propositional Formulas

A propositional formula φ is

- **satisfiable** if φ has at least one model
- **unsatisfiable** if φ is not satisfiable
- **valid** (or a **tautology**) if φ is true under every interpretation
- **falsifiable** if φ is no tautology

German: erfüllbar, unerfüllbar, allgemeingültig/eine Tautologie, falsifizierbar

Examples

How can we show that a formula has one of these properties?

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.

$\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)
- Show that $(A \wedge B)$ is **falsifiable**.

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.

$\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)

- Show that $(A \wedge B)$ is **falsifiable**.

$\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \not\models (A \wedge B)$)

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)
- Show that $(A \wedge B)$ is **falsifiable**.
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \not\models (A \wedge B)$)
- Show that $(A \wedge B)$ is **not valid**.

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)
- Show that $(A \wedge B)$ is **falsifiable**.
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \not\models (A \wedge B)$)
- Show that $(A \wedge B)$ is **not valid**.
Follows directly from falsifiability.

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)
- Show that $(A \wedge B)$ is **falsifiable**.
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \not\models (A \wedge B)$)
- Show that $(A \wedge B)$ is **not valid**.
Follows directly from falsifiability.
- Show that $(A \wedge B)$ is **not unsatisfiable**.

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)
- Show that $(A \wedge B)$ is **falsifiable**.
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \not\models (A \wedge B)$)
- Show that $(A \wedge B)$ is **not valid**.
Follows directly from falsifiability.
- Show that $(A \wedge B)$ is **not unsatisfiable**.
Follows directly from satisfiability.

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.

$\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)

- Show that $(A \wedge B)$ is **falsifiable**.

$\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \not\models (A \wedge B)$)

- Show that $(A \wedge B)$ is **not valid**.

Follows directly from falsifiability.

- Show that $(A \wedge B)$ is **not unsatisfiable**.

Follows directly from satisfiability.

So far all proofs by specifying **one** interpretation.

How to prove that a given formula is valid/unsatisfiable/
not satisfiable/not falsifiable?

Examples

How can we show that a formula has one of these properties?

- Show that $(A \wedge B)$ is **satisfiable**.
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \models (A \wedge B)$)
- Show that $(A \wedge B)$ is **falsifiable**.
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$ (+ simple proof that $\mathcal{I} \not\models (A \wedge B)$)
- Show that $(A \wedge B)$ is **not valid**.
Follows directly from falsifiability.
- Show that $(A \wedge B)$ is **not unsatisfiable**.
Follows directly from satisfiability.

So far all proofs by specifying **one** interpretation.

How to prove that a given formula is valid/unsatisfiable/
not satisfiable/not falsifiable?

\rightsquigarrow must consider **all possible** interpretations

Truth Tables

Evaluate for all possible interpretations
if they are models of the considered formula.

Truth Tables

Evaluate for all possible interpretations
if they are models of the considered formula.

$\mathcal{I}(A)$	$\mathcal{I} \models \neg A$
0	
1	

Truth Tables

Evaluate for all possible interpretations
if they are models of the considered formula.

$\mathcal{I}(A)$	$\mathcal{I} \models \neg A$
0	Yes
1	No

Truth Tables

Evaluate for all possible interpretations
if they are models of the considered formula.

$\mathcal{I}(A)$	$\mathcal{I} \models \neg A$
0	Yes
1	No

$\mathcal{I}(A)$	$\mathcal{I}(B)$	$\mathcal{I} \models (A \wedge B)$
0	0	
0	1	
1	0	
1	1	

Truth Tables

Evaluate for all possible interpretations
if they are models of the considered formula.

$\mathcal{I}(A)$	$\mathcal{I} \models \neg A$
0	Yes
1	No

$\mathcal{I}(A)$	$\mathcal{I}(B)$	$\mathcal{I} \models (A \wedge B)$
0	0	No
0	1	No
1	0	No
1	1	Yes

Truth Tables

Evaluate for all possible interpretations
if they are models of the considered formula.

$\mathcal{I}(A)$	$\mathcal{I} \models \neg A$
0	Yes
1	No

$\mathcal{I}(A)$	$\mathcal{I}(B)$	$\mathcal{I} \models (A \wedge B)$	$\mathcal{I}(A)$	$\mathcal{I}(B)$	$\mathcal{I} \models (A \vee B)$
0	0	No	0	0	No
0	1	No	0	1	Yes
1	0	No	1	0	Yes
1	1	Yes	1	1	Yes

Truth Tables in General

Similarly in the case where we consider a formula whose building blocks are themselves arbitrary unspecified formulas:

$\mathcal{I} \models \varphi$	$\mathcal{I} \models \psi$	$\mathcal{I} \models (\varphi \wedge \psi)$
No	No	No
No	Yes	No
Yes	No	No
Yes	Yes	Yes

Truth Tables for Properties of Formulas

Is $\varphi = ((A \rightarrow B) \vee (\neg B \rightarrow A))$ valid, unsatisfiable, ...?

$\mathcal{I}(A)$	$\mathcal{I}(B)$	$\mathcal{I} \models \neg B$	$\mathcal{I} \models (A \rightarrow B)$	$\mathcal{I} \models (\neg B \rightarrow A)$	$\mathcal{I} \models \varphi$
0	0	Yes	Yes	No	Yes
0	1	No	Yes	Yes	Yes
1	0	Yes	No	Yes	Yes
1	1	No	Yes	Yes	Yes

Connection Between Formula Properties and Truth Tables

A propositional formula φ is

- **satisfiable** if φ has at least one model
 \rightsquigarrow result in at least one row is “Yes”
- **unsatisfiable** if φ is not satisfiable
 \rightsquigarrow result in all rows is “No”
- **valid** (or a **tautology**) if φ is true under every interpretation
 \rightsquigarrow result in all rows is “Yes”
- **falsifiable** if φ is no tautology
 \rightsquigarrow result in at least one row is “No”

Main Disadvantage of Truth Tables

How big is a truth table with n atomic propositions?

Main Disadvantage of Truth Tables

How big is a truth table with n atomic propositions?

1		2 interpretations (rows)
2		4 interpretations (rows)
3		8 interpretations (rows)
n		??? interpretations

Main Disadvantage of Truth Tables

How big is a truth table with n atomic propositions?

1		2 interpretations (rows)
2		4 interpretations (rows)
3		8 interpretations (rows)
n		2^n interpretations

Some examples: $2^{10} = 1024$, $2^{20} = 1048576$, $2^{30} = 1073741824$

Main Disadvantage of Truth Tables

How big is a truth table with n atomic propositions?

1		2 interpretations (rows)
2		4 interpretations (rows)
3		8 interpretations (rows)
n		2^n interpretations

Some examples: $2^{10} = 1024$, $2^{20} = 1048576$, $2^{30} = 1073741824$

↪ not viable for larger formulas; we need a different solution

- more on difficulty of satisfiability etc.:
Theory of Computer Science course
- practical algorithms: Foundations of AI course

Discrete Mathematics in Computer Science

Equivalences

Malte Helmert, Gabriele Röger

University of Basel

Equivalent Formulas

Definition (Equivalence of Propositional Formulas)

Two propositional formulas φ and ψ over A are **(logically) equivalent** ($\varphi \equiv \psi$) if for **all interpretations \mathcal{I}** for A it is true that **$\mathcal{I} \models \varphi$ if and only if $\mathcal{I} \models \psi$** .

German: logisch äquivalent

Equivalent Formulas: Example

$$((\varphi \vee \psi) \vee \chi) \equiv (\varphi \vee (\psi \vee \chi))$$

$\mathcal{I} \models$ φ	$\mathcal{I} \models$ ψ	$\mathcal{I} \models$ χ	$\mathcal{I} \models$ $(\varphi \vee \psi)$	$\mathcal{I} \models$ $(\psi \vee \chi)$	$\mathcal{I} \models$ $((\varphi \vee \psi) \vee \chi)$	$\mathcal{I} \models$ $(\varphi \vee (\psi \vee \chi))$
No	No	No	No	No	No	No
No	No	Yes	No	Yes	Yes	Yes
No	Yes	No	Yes	Yes	Yes	Yes
No	Yes	Yes	Yes	Yes	Yes	Yes
Yes	No	No	Yes	No	Yes	Yes
Yes	No	Yes	Yes	Yes	Yes	Yes
Yes	Yes	No	Yes	Yes	Yes	Yes
Yes	Yes	Yes	Yes	Yes	Yes	Yes

Some Equivalences (1)

$$(\varphi \wedge \varphi) \equiv \varphi$$

$$(\varphi \vee \varphi) \equiv \varphi$$

(idempotence)

German: Idempotenz

Some Equivalences (1)

$$(\varphi \wedge \varphi) \equiv \varphi$$

$$(\varphi \vee \varphi) \equiv \varphi$$

(idempotence)

$$(\varphi \wedge \psi) \equiv (\psi \wedge \varphi)$$

$$(\varphi \vee \psi) \equiv (\psi \vee \varphi)$$

(commutativity)

German: Idempotenz, Kommutativität

Some Equivalences (1)

$$(\varphi \wedge \varphi) \equiv \varphi$$

$$(\varphi \vee \varphi) \equiv \varphi$$

(idempotence)

$$(\varphi \wedge \psi) \equiv (\psi \wedge \varphi)$$

$$(\varphi \vee \psi) \equiv (\psi \vee \varphi)$$

(commutativity)

$$((\varphi \wedge \psi) \wedge \chi) \equiv (\varphi \wedge (\psi \wedge \chi))$$

$$((\varphi \vee \psi) \vee \chi) \equiv (\varphi \vee (\psi \vee \chi))$$

(associativity)

German: Idempotenz, Kommutativität, Assoziativität

Some Equivalences (2)

$$(\varphi \wedge (\varphi \vee \psi)) \equiv \varphi$$

$$(\varphi \vee (\varphi \wedge \psi)) \equiv \varphi$$

(absorption)

German: Absorption

Some Equivalences (2)

$$(\varphi \wedge (\varphi \vee \psi)) \equiv \varphi$$

$$(\varphi \vee (\varphi \wedge \psi)) \equiv \varphi$$

(absorption)

$$(\varphi \wedge (\psi \vee \chi)) \equiv ((\varphi \wedge \psi) \vee (\varphi \wedge \chi))$$

$$(\varphi \vee (\psi \wedge \chi)) \equiv ((\varphi \vee \psi) \wedge (\varphi \vee \chi))$$

(distributivity)

German: Absorption, Distributivität

Some Equivalences (3)

$$\neg\neg\varphi \equiv \varphi$$

(double negation)

German: Doppelnegation

Some Equivalences (3)

$$\neg\neg\varphi \equiv \varphi$$

(double negation)

$$\neg(\varphi \wedge \psi) \equiv (\neg\varphi \vee \neg\psi)$$

$$\neg(\varphi \vee \psi) \equiv (\neg\varphi \wedge \neg\psi)$$

(De Morgan's rules)

German: Doppelnegation, De Morgansche Regeln

Some Equivalences (3)

$\neg\neg\varphi \equiv \varphi$ (double negation)

$\neg(\varphi \wedge \psi) \equiv (\neg\varphi \vee \neg\psi)$

$\neg(\varphi \vee \psi) \equiv (\neg\varphi \wedge \neg\psi)$ (De Morgan's rules)

$(\varphi \vee \psi) \equiv \varphi$ if φ tautology

$(\varphi \wedge \psi) \equiv \psi$ if φ tautology (tautology rules)

German: Doppelnegation, De Morgansche Regeln,
Tautologieregeln

Some Equivalences (3)

$\neg\neg\varphi \equiv \varphi$ (double negation)

$\neg(\varphi \wedge \psi) \equiv (\neg\varphi \vee \neg\psi)$

$\neg(\varphi \vee \psi) \equiv (\neg\varphi \wedge \neg\psi)$ (De Morgan's rules)

$(\varphi \vee \psi) \equiv \varphi$ if φ tautology

$(\varphi \wedge \psi) \equiv \psi$ if φ tautology (tautology rules)

$(\varphi \vee \psi) \equiv \psi$ if φ unsatisfiable

$(\varphi \wedge \psi) \equiv \varphi$ if φ unsatisfiable (unsatisfiability rules)

German: Doppelnegation, De Morgansche Regeln,
Tautologieregeln, Unerfüllbarkeitsregeln

Substitution Theorem

Theorem (Substitution Theorem)

Let φ and φ' be *equivalent* propositional formulas over A .

Let ψ be a propositional formula with (at least) one occurrence of the subformula φ .

Then ψ is *equivalent to* ψ' , where ψ' is constructed from ψ by *replacing* an occurrence of φ in ψ with φ' .

German: Ersetzbarkeitstheorem

(without proof)

Application of Equivalences: Example

$$(P \wedge (Q \vee \neg P)) \equiv ((P \wedge Q) \vee (P \wedge \neg P)) \quad (\text{distributivity})$$

Application of Equivalences: Example

$$\begin{aligned}(P \wedge (Q \vee \neg P)) &\equiv ((P \wedge Q) \vee (P \wedge \neg P)) && \text{(distributivity)} \\ &\equiv ((P \wedge \neg P) \vee (P \wedge Q)) && \text{(commutativity)}\end{aligned}$$

Application of Equivalences: Example

$$\begin{aligned}(P \wedge (Q \vee \neg P)) &\equiv ((P \wedge Q) \vee (P \wedge \neg P)) && \text{(distributivity)} \\ &\equiv ((P \wedge \neg P) \vee (P \wedge Q)) && \text{(commutativity)} \\ &\equiv (P \wedge Q) && \text{(unsatisfiability rule)}\end{aligned}$$