

Discrete Mathematics in Computer Science

A5. Proof Techniques II

Malte Helmert, Gabriele Röger

University of Basel

September 30, 2024

Discrete Mathematics in Computer Science

September 30, 2024 — A5. Proof Techniques II

A5.1 Mathematical Induction

A5.2 Structural Induction

A5.3 Excursus: Computer-assisted Theorem Proving

A5.1 Mathematical Induction

Proof Techniques

most common proof techniques:

- ▶ direct proof
- ▶ indirect proof (proof by contradiction)
- ▶ contrapositive
- ▶ **mathematical induction**
- ▶ structural induction

Mathematical Induction

Concrete Mathematics by Graham, Knuth and Patashnik (p. 3)
 Mathematical induction proves that
 we can climb as high as we like on a ladder,
 by proving that we can climb onto the bottom rung (the basis)
 and that
 from each rung we can climb up to the next one (the step).

Propositions

Consider a statement on all natural numbers n with $n \geq m$.

- ▶ E.g. "Every natural number $n \geq 2$ can be written as a product of prime numbers."
 - ▶ $P(2)$: "2 can be written as a product of prime numbers."
 - ▶ $P(3)$: "3 can be written as a product of prime numbers."
 - ▶ $P(4)$: "4 can be written as a product of prime numbers."
 - ▶ ...
 - ▶ $P(n)$: " n can be written as a product of prime numbers."
 - ▶ For every natural number $n \geq 2$ proposition $P(n)$ is true.

Proposition $P(n)$ is a mathematical statement that is defined in terms of natural number n .

Mathematical Induction

Mathematical Induction

Proof (of the truth) of proposition $P(n)$
 for all natural numbers n with $n \geq m$:

- ▶ **basis**: proof of $P(m)$
- ▶ **induction hypothesis** (IH):
 suppose that $P(k)$ is true for all k with $m \leq k \leq n$
- ▶ **inductive step**: proof of $P(n+1)$
 using the induction hypothesis

German: Vollständige Induktion, Induktionsanfang,
 Induktionsannahme oder Induktionsvoraussetzung,
 Induktionsschritt

Mathematical Induction: Example I

Theorem

Every natural number $n \geq 2$ can be written as a product of prime numbers, i. e. $n = p_1 \cdot p_2 \cdot \dots \cdot p_m$ with prime numbers p_1, \dots, p_m .

Proof.

Mathematical Induction over n :

basis $n = 2$: trivially satisfied, since 2 is prime

IH: Every natural number k with $2 \leq k \leq n$
 can be written as a product of prime numbers. ...

Mathematical Induction: Example I

Theorem

Every natural number $n \geq 2$ can be written as a product of prime numbers, i. e. $n = p_1 \cdot p_2 \cdot \dots \cdot p_m$ with prime numbers p_1, \dots, p_m .

Proof (continued).

inductive step $n \rightarrow n + 1$:

- ▶ **Case 1:** $n + 1$ is a prime number \rightsquigarrow trivial
- ▶ **Case 2:** $n + 1$ is not a prime number.
There are natural numbers $2 \leq q, r \leq n$ with $n + 1 = q \cdot r$.
Using the IH shows that there are prime numbers q_1, \dots, q_s with $q = q_1 \cdot \dots \cdot q_s$ and r_1, \dots, r_t with $r = r_1 \cdot \dots \cdot r_t$.
Together this means $n + 1 = q_1 \cdot \dots \cdot q_s \cdot r_1 \cdot \dots \cdot r_t$.

□

Mathematical Induction: Example II

Theorem

Let S be a finite set. Then $|\mathcal{P}(S)| = 2^{|S|}$.

What proposition can we use to prove this with mathematical induction?

Proof by Induction

Proof.

By induction over $|S|$.

Basis ($|S| = 0$): Then $S = \emptyset$ and $|\mathcal{P}(S)| = |\{\emptyset\}| = 1 = 2^0$.

IH: For all sets S with $|S| \leq n$, it holds that $|\mathcal{P}(S)| = 2^{|S|}$.

Inductive Step ($n \rightarrow n + 1$):

Let S' be an arbitrary set with $|S'| = n + 1$ and let e be an arbitrary member of S' .

Let further $S = S' \setminus \{e\}$ and $X = \{S'' \cup \{e\} \mid S'' \in \mathcal{P}(S)\}$.

Then $\mathcal{P}(S') = \mathcal{P}(S) \cup X$. As $\mathcal{P}(S)$ and X are disjoint and $|X| = |\mathcal{P}(S)|$, it holds that $|\mathcal{P}(S')| = 2|\mathcal{P}(S)|$.

Since $|S| = n$, we can use the IH and get

$$|\mathcal{P}(S')| = 2 \cdot 2^{|S|} = 2 \cdot 2^n = 2^{n+1} = 2^{|S'|}.$$

□

Weak vs. Strong Induction

- ▶ **Weak induction:** Induction hypothesis only supposes that $P(k)$ is true for $k = n$
- ▶ **Strong induction:** Induction hypothesis supposes that $P(k)$ is true for all $k \in \mathbb{N}_0$ with $m \leq k \leq n$
 - ▶ also: **complete induction**

Our previous definition corresponds to **strong induction**.

Which of the examples had also worked with weak induction?

Is Strong Induction More Powerful than Weak Induction?

Are there statements that we can prove with strong induction but not with weak induction?

We can always use a stronger proposition:

- ▶ “Every $n \in \mathbb{N}_0$ with $n \geq 2$ can be written as a product of prime numbers.”
- ▶ $P(n)$: “ n can be written as a product of prime numbers.”
- ▶ $P'(n)$: “all $k \in \mathbb{N}_0$ with $2 \leq k \leq n$ can be written as a product of prime numbers.”

A5.2 Structural Induction

Inductively Defined Sets: Examples

Example (Natural Numbers)

The set \mathbb{N}_0 of natural numbers is inductively defined as follows:

- ▶ 0 is a natural number.
- ▶ If n is a natural number, then $n + 1$ is a natural number.

Example (Binary Tree)

The set \mathcal{B} of binary trees is inductively defined as follows:

- ▶ \square is a binary tree (a leaf)
- ▶ If L and R are binary trees, then $\langle L, \circlearrowright, R \rangle$ is a binary tree (with inner node \circlearrowright).

Implicit statement: all elements of the set can be constructed by finite application of these rules

German: Binärbaum, Blatt, innerer Knoten

Inductive Definition of a Set

Inductive Definition

A set M can be defined **inductively** by specifying

- ▶ **basic elements** that are contained in M
- ▶ **construction rules** of the form
“Given some elements of M , another element of M can be constructed like this.”

German: Induktive Definition, Basiselemente, Konstruktionsregeln

Structural Induction

Structural Induction

Proof of statement for all elements of an inductively defined set

- ▶ **basis**: proof of the statement for the basic elements
- ▶ **induction hypothesis (IH)**:
suppose that the statement is true for some elements M
- ▶ **inductive step**: proof of the statement for elements constructed by applying a construction rule to M (one inductive step for each construction rule)

German: Strukturelle Induktion

Structural Induction: Example (1)

Definition (Leaves of a Binary Tree)

The number of **leaves** of a binary tree B , written $leaves(B)$, is defined as follows:

$$\begin{aligned} leaves(\square) &= 1 \\ leaves(\langle L, \circlearrowleft, R \rangle) &= leaves(L) + leaves(R) \end{aligned}$$

Definition (Inner Nodes of a Binary Tree)

The number of **inner nodes** of a binary tree B , written $inner(B)$, is defined as follows:

$$\begin{aligned} inner(\square) &= 0 \\ inner(\langle L, \circlearrowleft, R \rangle) &= inner(L) + inner(R) + 1 \end{aligned}$$

Structural Induction: Example (2)

Theorem

For all binary trees B : $inner(B) = leaves(B) - 1$.

Proof.

induction basis:

$$inner(\square) = 0 = 1 - 1 = leaves(\square) - 1$$

\rightsquigarrow statement is true for base case ...

Structural Induction: Example (3)

Proof (continued).

induction hypothesis:

to prove that the statement is true for a composite tree $\langle L, \circlearrowleft, R \rangle$, we may use that it is true for the subtrees L and R .

inductive step for $B = \langle L, \circlearrowleft, R \rangle$:

$$\begin{aligned} inner(B) &= inner(L) + inner(R) + 1 \\ &\stackrel{\text{IH}}{=} (leaves(L) - 1) + (leaves(R) - 1) + 1 \\ &= leaves(L) + leaves(R) - 1 = leaves(B) - 1 \end{aligned}$$

□

Structural Induction: Exercise

Definition (Height of a Binary Tree)

The **height** of a binary tree B , written $height(B)$, is defined as follows:

$$height(\square) = 0$$

$$height(\langle L, \circlearrowleft, R \rangle) = \max\{height(L), height(R)\} + 1$$

Prove by structural induction:

Theorem

For all binary trees B : $leaves(B) \leq 2^{height(B)}$.

Example: Tarradiddles

Example (Tarradiddles)

The set of tarradiddles is inductively defined as follows:

- ▶ \rightarrow is a tarradiddle.
- ▶ \heartsuit is a tarradiddle.
- ▶ If x and y are tarradiddles, then $x\clubsuit\clubsuit y$ is a tarradiddle.
- ▶ If x and y are tarradiddles, then $\clubsuit x \rightarrow y \clubsuit$ is a tarradiddle.

How do you prove with structural induction that every tarradiddle contains an even number of flowers?

A5.3 Excursus: Computer-assisted Theorem Proving


Computer-assisted Proofs

- ▶ Computers can help proving theorems.
- ▶ **Computer-aided proofs** have for example been used for proving theorems by exhaustion.
- ▶ Example: **Four color theorem**

Interactive Theorem Proving

- ▶ On the lowest abstraction level, rigorous mathematical proofs rely on formal logic.
- ▶ On this level, proofs can be automatically verified by computers.
- ▶ Nobody wants to write or read proofs on this level of detail.
- ▶ In Interactive Theorem Proving a human guides the proof and the computer tries to fill in the details.
- ▶ If it succeeds, we can be very confident that the proof is valid.
- ▶ Example theorem provers: Isabelle/HOL, Lean

Example



```

theory Mysets
  imports Main
begin

theorem set_example: "∀A,VB. (A-B = Set.empty ⟹ A⊆B)"
proof (rule ccontr)
  assume "¬(∀A,VB. (A-B = Set.empty ⟹ A⊆B))"
  hence "∃A,VB. (A-B = Set.empty ∧ ¬A⊆B)" by simp
  then obtain A::"a set" and B::"a set" where "A-B = Set.empty" "¬A⊆B" by simp
  hence "∃x. (x ∈ A ∧ x ∉ B)" by simp
  then obtain x::"a" where "(x ∈ A ∧ x ∉ B)" by (rule exE, simp)
  hence "x ∈ A-B" by simp
  hence "A-B ≠ Set.empty" using "x ∈ A-B" by simp
  with "A-B = Set.empty" show "False" by simp
qed
  
```

↔ Demo

Summary

- ▶ **Mathematical induction** is used to prove a proposition P for all natural numbers $\geq m$.
 - ▶ Prove $P(m)$.
 - ▶ Make hypothesis that $P(k)$ is true for $m \leq k \leq n$.
 - ▶ Establish $P(n+1)$ using the hypothesis.
- ▶ **Structural induction** applies the same general concept to prove a proposition P for all elements of an inductively defined set.