

Discrete Mathematics in Computer Science

B6. Sets: Comparing Cardinality and Hilbert's Hotel

Malte Helmert, Gabriele Röger

University of Basel

October 23, 2024

Comparing Cardinality

Finite Sets Revisited

We already know:

- The **cardinality** $|S|$ measures the size of set S .
- A set is **finite** if it has a finite number of elements.
- The **cardinality** of a finite set is the **number of elements** it contains.

Finite Sets Revisited

We already know:

- The **cardinality** $|S|$ measures the size of set S .
- A set is **finite** if it has a finite number of elements.
- The **cardinality** of a finite set is the **number of elements** it contains.

A set is **infinite** if it has an infinite number of elements.

Finite Sets Revisited

We already know:

- The **cardinality** $|S|$ measures the size of set S .
- A set is **finite** if it has a finite number of elements.
- The **cardinality** of a finite set is the **number of elements** it contains.

A set is **infinite** if it has an infinite number of elements.

Do all infinite sets have the same cardinality?

Comparing the Cardinality of Sets

- Consider $A = \{1, 2\}$ and $B = \{\text{dog}, \text{cat}, \text{mouse}\}$.
- We can map distinct elements of A to distinct elements of B , e.g.

$1 \mapsto \text{dog}$

$2 \mapsto \text{cat}$

Comparing the Cardinality of Sets

- Consider $A = \{1, 2\}$ and $B = \{\text{dog}, \text{cat}, \text{mouse}\}$.
- We can map distinct elements of A to distinct elements of B , e.g.

$1 \mapsto \text{dog}$

$2 \mapsto \text{cat}$

- This is an **injective function** from A to B :
 - every element of A is mapped to an element of B ;
 - different elements of A are mapped to different elements of B .

Comparing Cardinality

Definition (cardinality not larger)

Set A has **cardinality less than or equal** to the cardinality of set B ($|A| \leq |B|$), if **there is an injective function from A to B** .

Comparing the Cardinality of Sets

- $A = \{1, 2, 3\}$ and $B = \{\text{dog, cat, mouse}\}$ have cardinality 3.
- We can pair their elements by a bijection from A to B :

1 \leftrightarrow dog

2 \leftrightarrow cat

3 \leftrightarrow mouse

Comparing the Cardinality of Sets

- $A = \{1, 2, 3\}$ and $B = \{\text{dog, cat, mouse}\}$ have cardinality 3.
- We can pair their elements by a bijection from A to B :

1 \leftrightarrow dog

2 \leftrightarrow cat

3 \leftrightarrow mouse

- This is a **bijection** from A to B .
 - Each element of A is paired with exactly one element of set B .
 - Each element of B is paired with exactly one element of A .

Comparing the Cardinality of Sets

- $A = \{1, 2, 3\}$ and $B = \{\text{dog, cat, mouse}\}$ have cardinality 3.
- We can pair their elements by a bijection from A to B :

1 \leftrightarrow dog

2 \leftrightarrow cat

3 \leftrightarrow mouse

- This is a **bijection** from A to B .
 - Each element of A is paired with exactly one element of set B .
 - Each element of B is paired with exactly one element of A .

Comparing the Cardinality of Sets

- $A = \{1, 2, 3\}$ and $B = \{\text{dog, cat, mouse}\}$ have cardinality 3.
- We can pair their elements by a bijection from A to B :

1 \leftrightarrow dog

2 \leftrightarrow cat

3 \leftrightarrow mouse

- This is a **bijection** from A to B .
 - Each element of A is paired with exactly one element of set B .
 - Each element of B is paired with exactly one element of A .
- If there is a bijection from A to B there is one from B to A (the inverse function).

Equinumerous Sets

We use the existence of a bijection also as criterion for infinite sets:

Definition (equinumerous sets)

Two sets A and B have the same cardinality ($|A| = |B|$) if there **exists a bijection from A to B** .

Such sets are called **equinumerous**.

Equinumerous Sets

We use the existence of a bijection also as criterion for infinite sets:

Definition (equinumerous sets)

Two sets A and B have the same cardinality ($|A| = |B|$) if there **exists a bijection from A to B** .

Such sets are called **equinumerous**.

Definition (strictly smaller cardinality)

Set A has **cardinality strictly less** than the cardinality of set B ($|A| < |B|$), if $|A| \leq |B|$ and $|A| \neq |B|$.

Equinumerous Sets

We use the existence of a bijection also as criterion for infinite sets:

Definition (equinumerous sets)

Two sets A and B have the same cardinality ($|A| = |B|$) if there **exists a bijection from A to B** .

Such sets are called **equinumerous**.

Definition (strictly smaller cardinality)

Set A has **cardinality strictly less** than the cardinality of set B ($|A| < |B|$), if $|A| \leq |B|$ and $|A| \neq |B|$.

Consider set A and object $e \notin A$. Is $|A| < |A \cup \{e\}|$?

Questions



Questions?

Hilbert's Hotel


Hilbert's Hotel

Our intuition for finite sets does not always work for infinite sets.

Hilbert's Hotel

Our intuition for finite sets does not always work for infinite sets.


- If in a hotel all rooms are occupied then it cannot accommodate additional guests.



Hilbert's Hotel

Our intuition for finite sets does not always work for infinite sets.

- If in a hotel all rooms are occupied then it cannot accommodate additional guests.
- But **Hilbert's Grand Hotel** has **infinitely many rooms**.
- All these rooms are **occupied**.




One More Guest Arrives




- Every guest moves from her current room n to room $n + 1$.
- Room 1 is then free.
- The new guest gets room 1.

Four More Guests Arrive



- Every guest moves from her current room n to room $n + 4$.
- Rooms 1 to 4 are no longer occupied and can be used for the new guests.


Four More Guests Arrive




- Every guest moves from her current room n to room $n + 4$.
- Rooms 1 to 4 are no longer occupied and can be used for the new guests.

→ Works for any finite number of additional guests.

An Infinite Number of Guests Arrives



An Infinite Number of Guests Arrives



- Every guest moves from her current room n to room $2n$.
- The infinitely many rooms with odd numbers are now available.
- The new guests fit into these rooms.

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests

... arrive?

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests
- infinitely many ferries, each with an infinite number of coaches, each with infinitely many guests

... arrive?

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests
- infinitely many ferries, each with an infinite number of coaches, each with infinitely many guests
- ...

... arrive?

Can we Go further?

What if ...

- infinitely many coaches, each with an infinite number of guests
- infinitely many ferries, each with an infinite number of coaches, each with infinitely many guests
- ...

... arrive?

There are strategies for all these situations as long as with “infinite” we mean “countably infinite” and there is a finite number of layers.

Questions



Questions?

Questions



Questions?

Summary

Summary

- Set A has cardinality less than or equal the cardinality of set B ($|A| \leq |B|$), if there is an injective function from A to B .
- Sets A and B have the same cardinality ($|A| = |B|$) if there exists a bijection from A to B .

Summary

- Set A has cardinality less than or equal the cardinality of set B ($|A| \leq |B|$), if there is an injective function from A to B .
- Sets A and B have the same cardinality ($|A| = |B|$) if there exists a bijection from A to B .
- Our intuition for finite sets does not always work for infinite sets.