

# Discrete Mathematics in Computer Science

## D2. Properties of Formulas and Equivalences

Malte Helmert, Gabriele Röger

University of Basel

November 25/27, 2024

# Properties of Propositional Formulas

# The Story So Far

- **propositional logic** based on atomic propositions
- **syntax**: which formulas are well-formed?
- **semantics**: when is a formula true?
- **interpretations**: important basis of semantics

## Reminder: Syntax of Propositional Logic

### Definition (Syntax of Propositional Logic)

Let  $A$  be a set of **atomic propositions**. The set of **propositional formulas** (over  $A$ ) is inductively defined as follows:

- Every **atom**  $a \in A$  is a propositional formula over  $A$ .
- If  $\varphi$  is a propositional formula over  $A$ , then so is its **negation**  $\neg\varphi$ .
- If  $\varphi$  and  $\psi$  are propositional formulas over  $A$ , then so is the **conjunction**  $(\varphi \wedge \psi)$ .
- If  $\varphi$  and  $\psi$  are propositional formulas over  $A$ , then so is the **disjunction**  $(\varphi \vee \psi)$ .

The **implication**  $(\varphi \rightarrow \psi)$  is an abbreviation for  $(\neg\varphi \vee \psi)$ .

The **biconditional**  $(\varphi \leftrightarrow \psi)$  is an abbrev. for  $((\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi))$ .

## Reminder: Semantics of Propositional Logic

### Definition (Semantics of Propositional Logic)

A **truth assignment** (or **interpretation**) for a set of atomic propositions  $A$  is a function  $\mathcal{I} : A \rightarrow \{0, 1\}$ .

A propositional **formula**  $\varphi$  (over  $A$ ) **holds under**  $\mathcal{I}$  (written as  $\mathcal{I} \models \varphi$ ) according to the following definition:

|   |     |  |                  |
|---|-----|--|------------------|
| $\mathcal{I} \models a$                     | iff | $\mathcal{I}(a) = 1$   | (for $a \in A$ ) |
| $\mathcal{I} \models \neg\varphi$           | iff | not $\mathcal{I} \models \varphi$                            |                  |
| $\mathcal{I} \models (\varphi \wedge \psi)$ | iff | $\mathcal{I} \models \varphi$ and $\mathcal{I} \models \psi$ |                  |
| $\mathcal{I} \models (\varphi \vee \psi)$   | iff | $\mathcal{I} \models \varphi$ or $\mathcal{I} \models \psi$  |                  |

# Properties of Propositional Formulas

A propositional formula  $\varphi$  is

- **satisfiable** if  $\varphi$  has at least one model
- **unsatisfiable** if  $\varphi$  is not satisfiable
- **valid** (or a **tautology**) if  $\varphi$  is true under every interpretation
- **falsifiable** if  $\varphi$  is no tautology

**German:** erfüllbar, unerfüllbar, allgemeingültig/eine Tautologie, falsifizierbar

## Examples

How can we show that a formula has one of these properties?

## Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.



## Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.

$\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )

## Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.  
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **falsifiable**.

# Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.

$\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )

- Show that  $(A \wedge B)$  is **falsifiable**.

$\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \not\models (A \wedge B)$ )

# Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.  
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **falsifiable**.  
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \not\models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **not valid**.

## Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.  
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **falsifiable**.  
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \not\models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **not valid**.  
Follows directly from falsifiability.

# Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.  
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **falsifiable**.  
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \not\models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **not valid**.  
Follows directly from falsifiability.
- Show that  $(A \wedge B)$  is **not unsatisfiable**.

# Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.  
 $\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **falsifiable**.  
 $\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \not\models (A \wedge B)$ )
- Show that  $(A \wedge B)$  is **not valid**.  
Follows directly from falsifiability.
- Show that  $(A \wedge B)$  is **not unsatisfiable**.  
Follows directly from satisfiability.

# Examples

How can we show that a formula has one of these properties?

- Show that  $(A \wedge B)$  is **satisfiable**.

$\mathcal{I} = \{A \mapsto 1, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \models (A \wedge B)$ )

- Show that  $(A \wedge B)$  is **falsifiable**.

$\mathcal{I} = \{A \mapsto 0, B \mapsto 1\}$  (+ simple proof that  $\mathcal{I} \not\models (A \wedge B)$ )

- Show that  $(A \wedge B)$  is **not valid**.

Follows directly from falsifiability.

- Show that  $(A \wedge B)$  is **not unsatisfiable**.

Follows directly from satisfiability.

So far all proofs by specifying **one** interpretation.

How to prove that a given formula is valid/unsatisfiable/  
not satisfiable/not falsifiable?

$\rightsquigarrow$  must consider **all possible** interpretations



# Truth Tables

Evaluate for all possible interpretations  
if they are models of the considered formula.

# Truth Tables

Evaluate for all possible interpretations  
if they are models of the considered formula.

| $\mathcal{I}(A)$ | $\mathcal{I} \models \neg A$ |
|------------------|------------------------------|
| 0                |                              |
| 1                |                              |

# Truth Tables

Evaluate for all possible interpretations  
if they are models of the considered formula.

| $\mathcal{I}(A)$ | $\mathcal{I} \models \neg A$ |
|------------------|------------------------------|
| 0                | Yes                          |
| 1                | No                           |

# Truth Tables

Evaluate for all possible interpretations  
if they are models of the considered formula.

| $\mathcal{I}(A)$ | $\mathcal{I} \models \neg A$ |
|------------------|------------------------------|
| 0                | Yes                          |
| 1                | No                           |

| $\mathcal{I}(A)$ | $\mathcal{I}(B)$ | $\mathcal{I} \models (A \wedge B)$ |
|------------------|------------------|------------------------------------|
| 0                | 0                |                                    |
| 0                | 1                |                                    |
| 1                | 0                |                                    |
| 1                | 1                |                                    |

# Truth Tables

Evaluate for all possible interpretations  
if they are models of the considered formula.

| $\mathcal{I}(A)$ | $\mathcal{I} \models \neg A$ |
|------------------|------------------------------|
| 0                | Yes                          |
| 1                | No                           |

| $\mathcal{I}(A)$ | $\mathcal{I}(B)$ | $\mathcal{I} \models (A \wedge B)$ |
|------------------|------------------|------------------------------------|
| 0                | 0                | No                                 |
| 0                | 1                | No                                 |
| 1                | 0                | No                                 |
| 1                | 1                | Yes                                |

# Truth Tables

Evaluate for all possible interpretations  
if they are models of the considered formula.

| $\mathcal{I}(A)$ | $\mathcal{I} \models \neg A$ |
|------------------|------------------------------|
| 0                | Yes                          |
| 1                | No                           |

| $\mathcal{I}(A)$ | $\mathcal{I}(B)$ | $\mathcal{I} \models (A \wedge B)$ | $\mathcal{I}(A)$ | $\mathcal{I}(B)$ | $\mathcal{I} \models (A \vee B)$ |
|------------------|------------------|------------------------------------|------------------|------------------|----------------------------------|
| 0                | 0                | No                                 | 0                | 0                | No                               |
| 0                | 1                | No                                 | 0                | 1                | Yes                              |
| 1                | 0                | No                                 | 1                | 0                | Yes                              |
| 1                | 1                | Yes                                | 1                | 1                | Yes                              |

## Truth Tables in General

Similarly in the case where we consider a formula whose building blocks are themselves arbitrary unspecified formulas:

| $\mathcal{I} \models \varphi$ | $\mathcal{I} \models \psi$ | $\mathcal{I} \models (\varphi \wedge \psi)$ |
|-------------------------------|----------------------------|---|
| No                            | No                         | No  |
| No                            | Yes                        | No  |
| Yes                           | No                         | No  |
| Yes                           | Yes                        | Yes   |

## Truth Tables for Properties of Formulas

Is  $\varphi = ((A \rightarrow B) \vee (\neg B \rightarrow A))$  valid, unsatisfiable, ...?

| $\mathcal{I}(A)$ | $\mathcal{I}(B)$ | $\mathcal{I} \models \neg B$ | $\mathcal{I} \models (A \rightarrow B)$ | $\mathcal{I} \models (\neg B \rightarrow A)$ | $\mathcal{I} \models \varphi$ |
|------------------|------------------|------------------------------|---|--|-------------------------------|
| 0                | 0                | Yes                          | Yes                                     | No   | Yes                           |
| 0                | 1                | No                           | Yes                                     | Yes  | Yes                           |
| 1                | 0                | Yes                          | No                                      | Yes  | Yes                           |
| 1                | 1                | No                           | Yes                                     | Yes  | Yes                           |



# Connection Between Formula Properties and Truth Tables

A propositional formula  $\varphi$  is

- **satisfiable** if  $\varphi$  has at least one model  
     $\rightsquigarrow$  result in at least one row is “Yes”
- **unsatisfiable** if  $\varphi$  is not satisfiable  
     $\rightsquigarrow$  result in all rows is “No”
- **valid** (or a **tautology**) if  $\varphi$  is true under every interpretation  
     $\rightsquigarrow$  result in all rows is “Yes”
- **falsifiable** if  $\varphi$  is no tautology  
     $\rightsquigarrow$  result in at least one row is “No”

# Main Disadvantage of Truth Tables

How big is a truth table with  $n$  atomic propositions?

|     |  |                          |
|-----|--|--------------------------|
| 1   |  | 2 interpretations (rows) |
| 2   |  | 4 interpretations (rows) |
| 3   |  | 8 interpretations (rows) |
| $n$ |  | ??? interpretations      |

# Main Disadvantage of Truth Tables

How big is a truth table with  $n$  atomic propositions?

|     |  |                          |
|-----|--|--------------------------|
| 1   |  | 2 interpretations (rows) |
| 2   |  | 4 interpretations (rows) |
| 3   |  | 8 interpretations (rows) |
| $n$ |  | $2^n$ interpretations    |

Some examples:  $2^{10} = 1024$ ,  $2^{20} = 1048576$ ,  $2^{30} = 1073741824$

↪ not viable for larger formulas; we need a different solution

- more on difficulty of satisfiability etc.:  
Theory of Computer Science course
- practical algorithms: Foundations of AI course

# Equivalences

# Equivalent Formulas

## Definition (Equivalence of Propositional Formulas)

Two propositional formulas  $\varphi$  and  $\psi$  over  $A$  are **(logically) equivalent** ( $\varphi \equiv \psi$ ) if for **all interpretations  $\mathcal{I}$**  for  $A$  it is true that  **$\mathcal{I} \models \varphi$  if and only if  $\mathcal{I} \models \psi$** .

**German:** logisch äquivalent

## Equivalent Formulas: Example

$$((\varphi \vee \psi) \vee \chi) \equiv (\varphi \vee (\psi \vee \chi))$$

| $\mathcal{I} \models$<br>$\varphi$ | $\mathcal{I} \models$<br>$\psi$ | $\mathcal{I} \models$<br>$\chi$ | $\mathcal{I} \models$<br>$(\varphi \vee \psi)$ | $\mathcal{I} \models$<br>$(\psi \vee \chi)$ | $\mathcal{I} \models$<br>$((\varphi \vee \psi) \vee \chi)$ | $\mathcal{I} \models$<br>$(\varphi \vee (\psi \vee \chi))$ |
|------------------------------------|---------------------------------|---------------------------------|--|---|--|--|
| No                                 | No                              | No                              | No   | No  | No   | No   |
| No                                 | No                              | Yes                             | No   | Yes   | Yes  | Yes  |
| No                                 | Yes                             | No                              | Yes  | Yes   | Yes  | Yes  |
| No                                 | Yes                             | Yes                             | Yes  | Yes   | Yes  | Yes  |
| Yes                                | No                              | No                              | Yes  | No  | Yes  | Yes  |
| Yes                                | No                              | Yes                             | Yes  | Yes   | Yes  | Yes  |
| Yes                                | Yes                             | No                              | Yes  | Yes   | Yes  | Yes  |
| Yes                                | Yes                             | Yes                             | Yes  | Yes   | Yes  | Yes  |

## Some Equivalences (1)

$$(\varphi \wedge \varphi) \equiv \varphi$$

$$(\varphi \vee \varphi) \equiv \varphi$$

(idempotence)

German: Idempotenz

## Some Equivalences (1)

$$(\varphi \wedge \varphi) \equiv \varphi$$

$$(\varphi \vee \varphi) \equiv \varphi$$

(idempotence)

$$(\varphi \wedge \psi) \equiv (\psi \wedge \varphi)$$

$$(\varphi \vee \psi) \equiv (\psi \vee \varphi)$$

(commutativity)

German: Idempotenz, Kommutativität



## Some Equivalences (1)

$$(\varphi \wedge \varphi) \equiv \varphi$$

$$(\varphi \vee \varphi) \equiv \varphi$$

(idempotence)

$$(\varphi \wedge \psi) \equiv (\psi \wedge \varphi)$$

$$(\varphi \vee \psi) \equiv (\psi \vee \varphi)$$

(commutativity)

$$((\varphi \wedge \psi) \wedge \chi) \equiv (\varphi \wedge (\psi \wedge \chi))$$

$$((\varphi \vee \psi) \vee \chi) \equiv (\varphi \vee (\psi \vee \chi))$$

(associativity)

**German:** Idempotenz, Kommutativität, Assoziativität

## Some Equivalences (2)

$$(\varphi \wedge (\varphi \vee \psi)) \equiv \varphi$$

$$(\varphi \vee (\varphi \wedge \psi)) \equiv \varphi$$

(absorption)

German: Absorption

## Some Equivalences (2)

$$(\varphi \wedge (\varphi \vee \psi)) \equiv \varphi$$

$$(\varphi \vee (\varphi \wedge \psi)) \equiv \varphi$$

(absorption)

$$(\varphi \wedge (\psi \vee \chi)) \equiv ((\varphi \wedge \psi) \vee (\varphi \wedge \chi))$$

$$(\varphi \vee (\psi \wedge \chi)) \equiv ((\varphi \vee \psi) \wedge (\varphi \vee \chi))$$

(distributivity)

**German:** Absorption, Distributivität

## Some Equivalences (3)

$$\neg\neg\varphi \equiv \varphi$$

(double negation)

German: Doppelnegation

## Some Equivalences (3)

$$\neg\neg\varphi \equiv \varphi$$

(double negation)

$$\neg(\varphi \wedge \psi) \equiv (\neg\varphi \vee \neg\psi)$$

$$\neg(\varphi \vee \psi) \equiv (\neg\varphi \wedge \neg\psi)$$

(De Morgan's rules)

**German:** Doppelnegation, De Morgansche Regeln

## Some Equivalences (3)

$\neg\neg\varphi \equiv \varphi$  (double negation)

$\neg(\varphi \wedge \psi) \equiv (\neg\varphi \vee \neg\psi)$

$\neg(\varphi \vee \psi) \equiv (\neg\varphi \wedge \neg\psi)$  (De Morgan's rules)

$(\varphi \vee \psi) \equiv \varphi$  if  $\varphi$  tautology

$(\varphi \wedge \psi) \equiv \psi$  if  $\varphi$  tautology (tautology rules)

**German:** Doppelnegation, De Morgansche Regeln,  
Tautologieregeln

## Some Equivalences (3)

$$\neg\neg\varphi \equiv \varphi \quad \text{(double negation)}$$

$$\neg(\varphi \wedge \psi) \equiv (\neg\varphi \vee \neg\psi)$$

$$\neg(\varphi \vee \psi) \equiv (\neg\varphi \wedge \neg\psi) \quad \text{(De Morgan's rules)}$$

$$(\varphi \vee \psi) \equiv \varphi \text{ if } \varphi \text{ tautology}$$

$$(\varphi \wedge \psi) \equiv \psi \text{ if } \varphi \text{ tautology} \quad \text{(tautology rules)}$$

$$(\varphi \vee \psi) \equiv \psi \text{ if } \varphi \text{ unsatisfiable}$$

$$(\varphi \wedge \psi) \equiv \varphi \text{ if } \varphi \text{ unsatisfiable} \quad \text{(unsatisfiability rules)}$$

**German:** Doppelnegation, De Morgansche Regeln,  
Tautologieregeln, Unerfüllbarkeitsregeln

# Substitution Theorem

## Theorem (Substitution Theorem)

Let  $\varphi$  and  $\varphi'$  be *equivalent* propositional formulas over  $A$ .

Let  $\psi$  be a propositional formula with (at least) one occurrence of the subformula  $\varphi$ .

Then  $\psi$  is *equivalent to*  $\psi'$ , where  $\psi'$  is constructed from  $\psi$  by *replacing* an occurrence of  $\varphi$  in  $\psi$  with  $\varphi'$ .

German: Ersetzbarkeitstheorem

(without proof)



## Application of Equivalences: Example

$$(P \wedge (Q \vee \neg P)) \equiv ((P \wedge Q) \vee (P \wedge \neg P)) \quad (\text{distributivity})$$

## Application of Equivalences: Example

$$\begin{aligned}(P \wedge (Q \vee \neg P)) &\equiv ((P \wedge Q) \vee (P \wedge \neg P)) && \text{(distributivity)} \\ &\equiv ((P \wedge \neg P) \vee (P \wedge Q)) && \text{(commutativity)}\end{aligned}$$

## Application of Equivalences: Example

$$\begin{aligned}(P \wedge (Q \vee \neg P)) &\equiv ((P \wedge Q) \vee (P \wedge \neg P)) && \text{(distributivity)} \\ &\equiv ((P \wedge \neg P) \vee (P \wedge Q)) && \text{(commutativity)} \\ &\equiv (P \wedge Q) && \text{(unsatisfiability rule)}\end{aligned}$$