## Planning and Optimization


D2. Delete Relaxation: Properties of Relaxed Planning Tasks

Malte Helmert and Gabriele Röger

Universität Basel

October 21, 2024

#### Content of the Course



The Domination Lemma

## On-Set and Dominating States

#### Definition (On-Set)

The on-set of an interpretation s is the set of propositional variables that are true in s, i.e.,  $on(s) = s^{-1}(\{T\})$ .

→ for states of propositional planning tasks: states can be viewed as sets of (true) state variables

#### Definition (Dominate)

An interpretation s' dominates an interpretation s if  $on(s) \subseteq on(s')$ .

 $\rightsquigarrow$  all state variables true in s are also true in s'

## Domination Lemma (1)

#### Lemma (Domination)

Let s and s' be interpretations of a set of propositional variables V, and let  $\chi$  be a propositional formula over V which does not contain negation symbols.

If  $s \models \chi$  and s' dominates s, then  $s' \models \chi$ .

#### Proof.

Proof by induction over the structure of  $\chi$ .

- Base case  $\chi = \top$ : then  $s' \models \top$ .
- Base case  $\chi = \bot$ : then  $s \not\models \bot$ .

. . .

## Domination Lemma (2)

#### Proof (continued).

- Base case  $\chi = v \in V$ : if  $s \models v$ , then  $v \in on(s)$ . With  $on(s) \subseteq on(s')$ , we get  $v \in on(s')$  and hence  $s' \models v$ .
- Inductive case  $\chi = \chi_1 \wedge \chi_2$ : by induction hypothesis, our claim holds for the proper subformulas  $\chi_1$  and  $\chi_2$  of  $\chi$ .

■ Inductive case  $\chi = \chi_1 \vee \chi_2$ : analogous

## The Relaxation Lemma

#### Add Sets and Delete Sets

#### Definition (Add Set and Delete Set for an Effect)

Consider a propositional planning task with state variables V. Let e be an effect over V, and let s be a state over V. The add set of e in s, written addset(e, s), and the delete set of e in s, written delset(e, s), are defined as the following sets of state variables:

$$addset(e, s) = \{v \in V \mid s \models effcond(v, e)\}$$
$$delset(e, s) = \{v \in V \mid s \models effcond(\neg v, e)\}$$

Note: For all states s and operators o applicable in s, we have  $on(s[o]) = (on(s) \setminus delset(eff(o), s)) \cup addset(eff(o), s).$ 

#### Relaxation Lemma

For this and the following chapters on delete relaxation, we assume implicitly that we are working with propositional planning tasks in positive normal form.

#### Lemma (Relaxation)

Let s be a state, and let s' be a state that dominates s.

- If o is an operator applicable in s, then  $o^+$  is applicable in s' and  $s'[o^+]$  dominates s[o].
- 2 If  $\pi$  is an operator sequence applicable in s, then  $\pi^+$  is applicable in s' and s' $[\pi^+]$  dominates  $s[\pi]$ .
- **3** If additionally  $\pi$  leads to a goal state from state s, then  $\pi^+$  leads to a goal state from state s'.

## Proof of Relaxation Lemma (1)

#### Proof.

Let V be the set of state variables.

Part 1: Because o is applicable in s, we have  $s \models pre(o)$ .

Because pre(o) is negation-free and s' dominates s, we get  $s' \models pre(o)$  from the domination lemma.

Because  $pre(o^+) = pre(o)$ , this shows that  $o^+$  is applicable in s'.

. .

## Proof of Relaxation Lemma (2)

#### Proof (continued).

To prove that  $s'[o^+]$  dominates s[o], we first compare the relevant add sets:

$$addset(eff(o), s) = \{v \in V \mid s \models effcond(v, eff(o))\}$$

$$= \{v \in V \mid s \models effcond(v, eff(o^{+}))\} \qquad (1)$$

$$\subseteq \{v \in V \mid s' \models effcond(v, eff(o^{+}))\} \qquad (2)$$

$$= addset(eff(o^{+}), s'),$$

where (1) uses  $effcond(v, eff(o)) \equiv effcond(v, eff(o^+))$ and (2) uses the dominance lemma (note that effect conditions are negation-free for operators in positive normal form).

## Proof of Relaxation Lemma (3)

#### Proof (continued).

We then get:

$$on(s[o]) = (on(s) \setminus delset(eff(o), s)) \cup addset(eff(o), s)$$
  
 $\subseteq on(s) \cup addset(eff(o), s)$   
 $\subseteq on(s') \cup addset(eff(o^+), s')$   
 $= on(s'[o^+]),$ 

and thus  $s'[o^+]$  dominates s[o].

This concludes the proof of Part 1.

## Proof of Relaxation Lemma (4)

#### Proof (continued).

Part 2: by induction over  $n = |\pi|$ 

Base case:  $\pi = \langle \rangle$ 

The empty plan is trivially applicable in s', and  $s' \llbracket \langle \rangle^+ \rrbracket = s'$  dominates  $s \llbracket \langle \rangle \rrbracket = s$  by prerequisite.

Inductive case:  $\pi = \langle o_1, \dots, o_{n+1} \rangle$ 

By the induction hypothesis,  $\langle o_1^+, \ldots, o_n^+ \rangle$  is applicable in s', and  $t' = s' [\langle o_1^+, \dots, o_n^+ \rangle]$  dominates  $t = s [\langle o_1, \dots, o_n \rangle]$ .

Also,  $o_{n+1}$  is applicable in t.

Using Part 1,  $o_{n+1}^+$  is applicable in t' and  $s'[\pi^+] = t'[o_{n+1}^+]$ dominates  $s[\pi] = t[o_{n+1}]$ .

This concludes the proof of Part 2.

## Proof of Relaxation Lemma (5)

#### Proof (continued).

Part 3: Let  $\gamma$  be the goal formula.

From Part 2, we obtain that  $t' = s' \llbracket \pi^+ \rrbracket$  dominates  $t = s \llbracket \pi \rrbracket$ . By prerequisite, t is a goal state and hence  $t \models \gamma$ .

Because the task is in positive normal form,  $\gamma$  is negation-free, and hence  $t' \models \gamma$  because of the domination lemma.

Therefore, t' is a goal state.



## Consequences

### Consequences of the Relaxation Lemma

- The relaxation lemma is the main technical result that we will use to study delete relaxation.
- Next, we show two further properties of delete relaxation that will be useful for us.
- They are direct consequences of the relaxation lemma.

### Consequences of the Relaxation Lemma (1)

#### Corollary (Relaxation Preserves Plans and Leads to Dominance)

Let  $\pi$  be an operator sequence that is applicable in state s. Then  $\pi^+$  is applicable in s and  $s[\pi^+]$  dominates  $s[\pi]$ . If  $\pi$  is a plan for  $\Pi$ , then  $\pi^+$  is a plan for  $\Pi^+$ .

#### Proof.

Apply relaxation lemma with s' = s.

- Nelaxations of plans are relaxed plans.
- → Delete relaxation is no harder to solve than original task.
- Optimal relaxed plans are never more expensive than optimal plans for original tasks.

#### Corollary (Relaxation Preserves Dominance)

Let s be a state, let s' be a state that dominates s, and let  $\pi^+$  be a relaxed operator sequence applicable in s.

Then  $\pi^+$  is applicable in s' and s' $[\pi^+]$  dominates  $s[\pi^+]$ .

#### Proof.

Apply relaxation lemma with  $\pi^+$  for  $\pi$ , noting that  $(\pi^+)^+ = \pi^+$ .

- $\rightarrow$  If there is a relaxed plan starting from state s, the same plan can be used starting from a dominating state s'.
- → Dominating states are always "better" in relaxed tasks.


# Monotonicity

#### Lemma (Monotonicity)

Let s be a state in which relaxed operator  $o^+$  is applicable. Then  $s[o^+]$  dominates s.

#### Proof.

Since relaxed operators only have positive effects, we have  $on(s) \subseteq on(s) \cup addset(eff(o^+), s) = on(s[o^+])$ .



## Finding Relaxed Plans

Using the theory we developed, we are now ready to study the problem of finding plans for relaxed planning tasks.

→ next chapter

# Summary

### Summary

- With positive normal form, having more true variables is good.
- We can formalize this as dominance between states.
- It follows that delete relaxation is a simplification: it is never harder to solve a relaxed task than the original one.
- In delete-relaxed tasks, applying an operator always takes us to a dominating state and therefore never hurts.