

# Planning and Optimization

## F6. Linear & Integer Programming

Malte Helmert and Gabriele Röger

Universität Basel

December 4, 2024

# Planning and Optimization

December 4, 2024 — F6. Linear & Integer Programming


F6.1 Integer Programs

F6.2 Linear Programs


F6.3 Normal Forms and Duality

F6.4 Summary


## Content of the Course (Timeline)



## Content of the Course (Relevance)



## Not Content of this Course (Relevance)



## F6.1 Integer Programs

## Motivation

- ▶ This goes on beyond Computer Science
- ▶ Active **research** on IPs and LPs in
  - ▶ Operation Research
  - ▶ Mathematics
- ▶ Many **application** areas, for instance:
  - ▶ Manufacturing
  - ▶ Agriculture
  - ▶ Mining
  - ▶ Logistics
  - ▶ **Planning**
- ▶ As an application, we treat LPs / IPs as a **blackbox**
- ▶ We just look at **the fundamentals**
- ▶ However, even on the application side there is much more (e.g., modelling tricks or solver parameters to speed up computation)

## Motivation

### Example (Optimization Problem)

Consider the following scenario:

- ▶ A factory produces two products A and B
- ▶ Selling one (unit of) B yields 5 times the profit of selling one A
- ▶ A client places the unusual order to “buy anything that can be produced on that day as long as two plus twice the units of A is not smaller than the number of B”
- ▶ More than 12 products in total cannot be produced per day
- ▶ There is only material for 6 units of A (there is enough material to produce any amount of B)

How many units of A and B does the client receive if the factory owner aims to maximize her profit?

## Integer Program: Example

Let  $X_A$  and  $X_B$  be the (integer) number of produced A and B

Example (Optimization Problem as Integer Program)

maximize  $X_A + 5X_B$  subject to

$$2 + 2X_A \geq X_B$$

$$X_A + X_B \leq 12$$

$$X_A \leq 6$$

$$X_A \geq 0, \quad X_B \geq 0$$

↪ unique optimal solution:

produce 4 A ( $X_A = 4$ ) and 8 B ( $X_B = 8$ ) for a profit of 44

## Same Program as Input for the CPLEX Solver

File ip.lp

Maximize

$$\text{obj: } X\_A + 5 X\_B$$

Subject To

$$c1: -2 X\_A + X\_B \leq 2$$

$$c2: X\_A + X\_B \leq 12$$

Bounds

$$0 \leq X\_A \leq 6$$

$$0 \leq X\_B$$


General

$$X\_A \quad X\_B$$

End

→ Demo

## Integer Program Example: Visualization



## Integer Programs

Integer Program

An **integer program (IP)** consists of:

- ▶ a finite set of **integer-valued variables**  $V$
- ▶ a finite set of **linear inequalities** (constraints) over  $V$
- ▶ an **objective function**, which is a linear combination of  $V$
- ▶ which should be **minimized** or **maximized**.

## Terminology

- ▶ An integer assignment to all variables in  $V$  is **feasible** if it satisfies the constraints.
- ▶ An integer program is **feasible** if there is such a feasible assignment. Otherwise it is **infeasible**.
- ▶ A feasible maximum (resp. minimum) problem is **unbounded** if the objective function can assume arbitrarily large positive (resp. negative) values at feasible assignments. Otherwise it is **bounded**.
- ▶ The **objective value** of a bounded feasible maximum (resp. minimum) problem is the maximum (resp. minimum) value of the objective function with a feasible assignment.

## Another Example

### Example

minimize  $3X_{o_1} + 4X_{o_2} + 5X_{o_3}$  subject to

$$X_{o_4} \geq 1$$

$$X_{o_1} + X_{o_2} \geq 1$$

$$X_{o_1} + X_{o_3} \geq 1$$

$$X_{o_2} + X_{o_3} \geq 1$$

$$X_{o_1} \geq 0, \quad X_{o_2} \geq 0, \quad X_{o_3} \geq 0, \quad X_{o_4} \geq 0$$

What example from a recent chapter does this IP encode?

↪ the **minimum hitting set** from Chapter F4

## Complexity of Solving Integer Programs

- ▶ As an IP can compute an MHS, solving an IP must be **at least as complex** as computing an MHS
- ▶ Reminder: MHS is a “classical” NP-complete problem
- ▶ Good news: Solving an IP is **not harder**

↪ Finding solutions for IPs is **NP-complete**.

Removing the requirement that solutions must be **integer-valued** leads to a simpler problem

## F6.2 Linear Programs

## Linear Programs

### Linear Program

A **linear program (LP)** consists of:

- ▶ a finite set of **real-valued variables**  $V$
- ▶ a finite set of **linear inequalities** (constraints) over  $V$
- ▶ an **objective function**, which is a linear combination of  $V$
- ▶ which should be **minimized** or **maximized**.

We use the introduced IP terminology also for LPs.

**Mixed IPs (MIPs)** are something between IPs and LPs:  
some variables are integer-valued, some are real-valued.

## Linear Program: Example

Let  $X_A$  and  $X_B$  be the (**real-valued**) number of produced A and B

**Example (Optimization Problem as Linear Program)**

maximize  $X_A + 5X_B$  subject to

$$2 + 2X_A \geq X_B$$

$$X_A + X_B \leq 12$$

$$X_A \leq 6$$

$$X_A \geq 0, \quad X_B \geq 0$$

↪ unique optimal solution:

$$X_A = 3\frac{1}{3} \text{ and } X_B = 8\frac{2}{3} \text{ with objective value } 46\frac{2}{3}$$

## Same Program as Input for the CPLEX Solver

### File lp.lp

Maximize

$$\text{obj: } X_A + 5 X_B$$

Subject To

$$c1: -2 X_A + X_B \leq 2$$

$$c2: X_A + X_B \leq 12$$

Bounds


$$0 \leq X_A \leq 6$$

$$0 \leq X_B$$

End

→ Demo

## Linear Program Example: Visualization



## Solving Linear Programs

- ▶ **Observation:**  
Here, LP solution is an **upper bound** for the corresponding IP.
- ▶ **Complexity:**  
LP solving is a **polynomial-time** problem.
- ▶ **Common idea:**  
Approximate IP solution with corresponding LP (**LP relaxation**).

## LP Relaxation

### Theorem (LP Relaxation)

The **LP relaxation** of an integer program is the problem that arises by removing the requirement that variables are integer-valued.

For a **maximization** (resp. **minimization**) problem, the objective value of the LP relaxation is an **upper** (resp. **lower**) **bound** on the value of the IP.

### Proof idea.

Every feasible assignment for the IP is also feasible for the LP.  $\square$

## LP Relaxation of MHS heuristic

### Example (Minimum Hitting Set)

minimize  $3X_{o_1} + 4X_{o_2} + 5X_{o_3}$  subject to

$$\begin{aligned} X_{o_4} &\geq 1 \\ X_{o_1} + X_{o_2} &\geq 1 \\ X_{o_1} + X_{o_3} &\geq 1 \\ X_{o_2} + X_{o_3} &\geq 1 \end{aligned}$$

$$X_{o_1} \geq 0, \quad X_{o_2} \geq 0, \quad X_{o_3} \geq 0, \quad X_{o_4} \geq 0$$

- ↪ optimal solution of **LP relaxation**:  
 $X_{o_4} = 1$  and  $X_{o_1} = X_{o_2} = X_{o_3} = 0.5$  with objective value 6
- ↪ LP relaxation of MHS heuristic is **admissible**  
and can be computed in **polynomial time**

## F6.3 Normal Forms and Duality

## Standard Maximum Problem

Normal form for maximization problems:

**Definition (Standard Maximum Problem)**

Find values for  $x_1, \dots, x_n$ , to maximize

$$c_1x_1 + c_2x_2 + \dots + c_nx_n$$

subject to the constraints

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &\leq b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &\leq b_2 \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &\leq b_m \end{aligned}$$

and  $x_1 \geq 0, x_2 \geq 0, \dots, x_n \geq 0$ .

## Standard Maximum Problem: Matrix and Vectors

A standard maximum problem is often given by

- ▶ an  $m$ -vector  $\mathbf{b} = \langle b_1, \dots, b_m \rangle^T$  (**bounds**),
- ▶ an  $n$ -vector  $\mathbf{c} = \langle c_1, \dots, c_n \rangle^T$  (**objective coefficients**),
- ▶ and an  $m \times n$  matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix} \quad (\text{coefficients})$$

- ▶ Then the problem is to find a vector  $\mathbf{x} = \langle x_1, \dots, x_n \rangle^T$  to maximize  $\mathbf{c}^T \mathbf{x}$  subject to  $\mathbf{A}\mathbf{x} \leq \mathbf{b}$  and  $\mathbf{x} \geq \mathbf{0}$ .

## Standard Minimum Problem

- ▶ there is also a **standard minimum problem**
- ▶ it's form is identical to the standard maximum problem, except that
  - ▶ the aim is to minimize the objective function
  - ▶ subject to  $\mathbf{A}\mathbf{x} \geq \mathbf{b}$
- ▶ All linear programs can efficiently be converted into a standard maximum/minimum problem.

## Some LP Theory: Duality

Every LP has an alternative view (its **dual LP**).

| Primal                         | Dual                           |
|--------------------------------|--------------------------------|
| maximization (or minimization) | minimization (or maximization) |
| objective coefficients         | bounds                         |
| bounds                         | objective coefficients         |
| bounded variable               | $\geq$ -constraint             |
| $\leq$ -constraint             | bounded variable               |
| free variable                  | $=$ -constraint                |
| $=$ -constraint                | free variable                  |

dual of dual: original LP

## Dual Problem

### Definition (Dual Problem)

The **dual** of the standard maximum problem

$$\text{maximize } \mathbf{c}^T \mathbf{x} \text{ subject to } \mathbf{Ax} \leq \mathbf{b} \text{ and } \mathbf{x} \geq \mathbf{0}$$

is the standard minimum problem

$$\text{minimize } \mathbf{b}^T \mathbf{y} \text{ subject to } \mathbf{A}^T \mathbf{y} \geq \mathbf{c} \text{ and } \mathbf{y} \geq \mathbf{0}$$

## Dual Problem: Example

### Example (Dual of the Optimization Problem)

maximize  $X_A + 5X_B$  subject to

$$[Y_1] \quad -2X_A + X_B \leq 2$$

$$[Y_2] \quad X_A + X_B \leq 12$$

$$[Y_3] \quad X_A \leq 6$$

$$X_A \geq 0, \quad X_B \geq 0$$

## Dual Problem: Example

### Example (Dual of the Optimization Problem)

minimize  $2Y_1 + 12Y_2 + 6Y_3$  subject to

$$[X_A] \quad -2Y_1 + Y_2 + Y_3 \geq 1$$

$$[X_B] \quad Y_1 + Y_2 \geq 5$$

$$Y_1 \geq 0, \quad Y_2 \geq 0, \quad Y_3 \geq 0$$

## Duality Theorem

### Theorem (Duality Theorem)

If a standard linear program is **bounded feasible**, then so is its dual, and their **objective values are equal**.

(Proof omitted.)

The dual provides a different perspective on a problem.




## F6.4 Summary

## Summary

- ▶ **Linear (and integer) programs** consist of an **objective function** that should be **maximized or minimized** subject to a set of given **linear constraints**.
- ▶ Finding solutions for **integer** programs is **NP-complete**.
- ▶ **LP solving** is a **polynomial time** problem.
- ▶ The dual of a maximization LP is a minimization LP and vice versa.
- ▶ The **dual** of a bounded feasible LP has the **same objective value**.

## Further Reading

The slides in this chapter are based on the following excellent tutorial on LP solving:

-  **Thomas S. Ferguson.**  
Linear Programming – A Concise Introduction.  
UCLA, unpublished document available online.