

Discrete Mathematics in Computer Science

B4. Operations on Relations

Malte Helmert, Gabriele Röger

University of Basel

October 13, 2025

Discrete Mathematics in Computer Science

October 13, 2025 — B4. Operations on Relations

B4.1 Operations on Relations

B4.1 Operations on Relations

Relations: Recap

- ▶ A **relation over sets** $S_1, \dots, S_n$ is a set $R \subseteq S_1 \times \dots \times S_n$.
- ▶ A **binary** relation is a relation over two sets.
- ▶ A **homogeneous** relation R over set S is a binary relation $R \subseteq S \times S$.

Set Operations

- ▶ Relations are **sets** of tuples, so we can build their union, intersection, complement,
- ▶ Let R be a relation over $S_1, \dots, S_n$ and R' a relation over $S'_1, \dots, S'_n$. Then $R \cup R'$ is a relation over $S_1 \cup S'_1, \dots, S_n \cup S'_n$.
With the standard relations $<, =$ and $\leq$ for $\mathbb{N}_0$,
relation $\leq$ corresponds to the union of relations $<$ and $=$.
- ▶ Let R and R' be relations over n sets.
Then $R \cap R'$ is a relation.
Over which sets?
With the standard relations $\leq, =$ and $\geq$ for $\mathbb{N}_0$,
relation $=$ corresponds to the intersection of $\leq$ and $\geq$.
- ▶ If R is a relation over $S_1, \dots, S_n$
then so is the **complementary relation** $\bar{R} = (S_1 \times \dots \times S_n) \setminus R$.
With the standard relations for $\mathbb{N}_0$, relation $=$ is the
complementary relation of $\neq$ and $>$ the one of $\leq$.

Inverse of a Relation

Definition

Let $R \subseteq A \times B$ be a binary relation over A and B .

The **inverse relation** of R is the relation $R^{-1} \subseteq B \times A$ given by $R^{-1} = \{(b, a) \mid (a, b) \in R\}$.

- ▶ The inverse of the $<$ relation over $\mathbb{N}_0$ is the $>$ relation.
- ▶ Relation R with xRy iff person x has a key for y .
Inverse: Q with aQb iff lock a can be opened by person b .

German: inverse Relation oder Umkehrrelation

Composition of Relations

Definition (Composition of relations)

Let R_1 be a relation over A and B and R_2 a relation over B and C . The **composition of R_1 and R_2** is the relation $R_2 \circ R_1$ over A and C with:

$$R_2 \circ R_1 = \{(a, c) \mid \text{there is a } b \in B \text{ with} \\ (a, b) \in R_1 \text{ and } (b, c) \in R_2\}$$

How can we illustrate this graphically?

German: Komposition oder Rückwärtsverkettung

Composition of Relations: Example

$$S_1 = \{1, 2, 3, 4\}$$

$$S_2 = \{A, B, C, D, E\}$$

$$S_3 = \{a, b, c, d\}$$

$$R_1 = \{(1, A), (1, B), (3, B), (4, D)\} \text{ over } S_1 \text{ and } S_2$$

$$R_2 = \{(B, a), (C, c), (D, a), (D, d)\} \text{ over } S_2 \text{ and } S_3$$

$$R_2 \circ R_1 =$$

Composition is Associative

Theorem (Associativity of composition)

Let $S_1, \dots, S_4$ be sets and R_1, R_2, R_3 relations with $R_i \subseteq S_i \times S_{i+1}$.
Then

$$R_3 \circ (R_2 \circ R_1) = (R_3 \circ R_2) \circ R_1.$$

Proof.

It holds that $(x_1, x_4) \in R_3 \circ (R_2 \circ R_1)$ iff there is an x_3 with $(x_1, x_3) \in R_2 \circ R_1$ and $(x_3, x_4) \in R_3$.

As $(x_1, x_3) \in R_2 \circ R_1$ iff there is an x_2 with $(x_1, x_2) \in R_1$ and $(x_2, x_3) \in R_2$, we have overall that $(x_1, x_4) \in R_3 \circ (R_2 \circ R_1)$ iff there are x_2, x_3 with $(x_1, x_2) \in R_1$, $(x_2, x_3) \in R_2$ and $(x_3, x_4) \in R_3$.

This is the case iff there is an x_2 with $(x_1, x_2) \in R_1$ and $(x_2, x_4) \in R_3 \circ R_2$, which holds iff $(x_1, x_4) \in (R_3 \circ R_2) \circ R_1$. □

(Reflexive) Transitive Closure

Definition ((Reflexive) transitive closure)

Let R be a relation over set S .

The **transitive closure** R^+ of R is the **smallest relation over S that is transitive and has R as a subset.**

The **reflexive transitive closure** R^* of R is the **smallest relation over S that is reflexive, transitive and has R as a subset.**

The (reflexive) transitive closure always exists. **Why?**

Example: If aRb specifies that there is a direct flight from a to b , what do R^+ and R^* express?

German: (reflexive) transitive Hülle

Transitive Closure and n -fold Composition

Define the n -fold composition of a relation R over S as

$$\begin{aligned} R_0 &= \{(x, x) \mid x \in S\} & \text{and} \\ R_i &= R \circ R_{i-1} & \text{for } i \geq 1. \end{aligned}$$

Theorem

Let R be a relation over set S .

Then $R^+ = \bigcup_{i=1}^{\infty} R_i$ and $R^* = \bigcup_{i=0}^{\infty} R_i$.

Without proof.

German: n -fache Komposition

Other Operators

- ▶ There are many more operators, also for general relations.
- ▶ Highly relevant for **queries over relational databases**.
- ▶ For example, **join** operators combine relations based on common entries.
- ▶ Example for a **natural join**:

Employee

Name	Empld	DeptName
Harry	3415	Finance
Sally	2241	Sales
George	3401	Finance
Harriet	2202	Sales
Mary	1257	Human Resources

Dept

DeptName	Manager
Finance	George
Sales	Harriet
Production	Charles

Employee ⋈ Dept

Name	Empld	DeptName	Manager
Harry	3415	Finance	George
Sally	2241	Sales	Harriet
George	3401	Finance	George
Harriet	2202	Sales	Harriet

(Source: Wikipedia)

Summary

- ▶ Relations: general, binary, homogeneous
- ▶ Properties: reflexivity, symmetry, transitivity
(and related properties)
- ▶ Special relations: equivalence relations, order relations
- ▶ Operations: inverse, composition, transitive closure