Planning and Optimization

C5. SAT Planning: Core Idea and Sequential Encoding

Malte Helmert and Gabriele Röger

Universität Basel

October 12, 2026

C5.1 Introduction

C5.2 Formula Overview

C5.3 Initial State, Goal, Operator Selection

C5.4 Transitions

C5.5 Summary

Content of the Course



C5.1 Introduction

SAT Solvers

- SAT solvers (algorithms that find satisfying assignments to CNF formulas) are one of the major success stories in solving hard combinatorial problems.
- Can we leverage them for classical planning?

$\rightsquigarrow$ SAT planning (a.k.a. planning as satisfiability)

background on SAT Solvers:
$\rightsquigarrow$ Foundations of Artificial Intelligence Course, Ch. E4–E5

Complexity Mismatch

- The SAT problem is NP-complete, while PlanEx is PSPACE-complete.

$\rightsquigarrow$ one-shot polynomial reduction from PlanEx to SAT not possible (unless NP = PSPACE)

Solution: Iterative Deepening

- We can generate a propositional formula that tests if task Π has a plan with horizon (length bound) T in time $O(\|\Pi\|^k \cdot T)$ ($\rightsquigarrow$ pseudo-polynomial reduction).
- Use as building block of algorithm that probes increasing horizons (a bit like IDA*).
- Can be efficient if there exist plans that are not excessively long.

SAT Planning: Main Loop

basic SAT planning algorithm:

SAT Planning

```
def satplan(Π):
    for T ∈ {0, 1, 2, ...}:
        φ := build_sat_formula(Π, T)
        I = sat_solver(φ)                ▷ returns a model or none
        if I is not none:
            return extract_plan(Π, T, I)
```

Termination criterion for unsolvable tasks?

C5.2 Formula Overview

SAT Formula: CNF?

- SAT solvers require conjunctive normal form (CNF), i.e., formulas expressed as collection of clauses.
- We will make sure that our SAT formulas are in CNF when our input is a STRIPS task.
- We do allow fully general propositional tasks, but then the formula may need additional conversion to CNF.

SAT Formula: Variables

- given propositional planning task $\Pi = \langle V, I, O, \gamma \rangle$
- given horizon $T \in \mathbb{N}_0$

Variables of the SAT Formula

- propositional variables v^i for all $v \in V$, $0 \leq i \leq T$ encode state after i steps of the plan
- propositional variables o^i for all $o \in O$, $1 \leq i \leq T$ encode operator(s) applied in i-th step of the plan

Formulas with Time Steps

Definition (Time-Stamped Formulas)

Let φ be a propositional logic formula over the variables V.
Let $0 \leq i \leq T$.

We write φ^i for the formula obtained from φ by replacing each $v \in V$ with v^i.

Example: $((a \wedge b) \vee \neg c)^3 = (a^3 \wedge b^3) \vee \neg c^3$

SAT Formula: Motivation

We want to express a formula whose models are exactly the plans/traces with T steps.

For this, the formula must express four things:

- The variables v^0 ($v \in V$) define the initial state.
- The variables v^T ($v \in V$) define a goal state.
- We select exactly one operator variable o^i ($o \in O$) for each time step $1 \leq i \leq T$.
- If we select o^i, then variables v^{i-1} and v^i ($v \in V$) describe a state transition from the $(i-1)$-th state of the plan to the i-th state of the plan (that uses operator o).

The final formula is the conjunction of all these parts.

C5.3 Initial State, Goal, Operator Selection

SAT Formula: Initial State

SAT Formula: Initial State

initial state clauses:

- v^0 for all $v \in V$ with $I(v) = \mathbf{T}$
- $\neg v^0$ for all $v \in V$ with $I(v) = \mathbf{F}$

SAT Formula: Goal

SAT Formula: Goal

goal clauses:

- γ^T

For STRIPS, this is a conjunction of unit clauses.
For general goals, this may not be in clause form.

SAT Formula: Operator Selection

Let $O = \{o_1, \dots, o_n\}$.

SAT Formula: Operator Selection

operator selection clauses:

- $o_1^i \vee \cdots \vee o_n^i$ for all $1 \leq i \leq T$

operator exclusion clauses:

- $\neg o_j^i \vee \neg o_k^i$ for all $1 \leq i \leq T$, $1 \leq j < k \leq n$

C5.4 Transitions

SAT Formula: Transitions

We now get to the interesting/challenging bit:
encoding the transitions.

Key observations: if we apply operator o at time i,

- its precondition must be satisfied at time $i-1$:
  $o^i \rightarrow pre(o)^{i-1}$
- variable v is true at time i iff its regression is true at $i-1$:
  $o^i \rightarrow (v^i \leftrightarrow regr(v, eff(o))^{i-1})$

Question: Why $regr(v, eff(o))$, not $regr(v, o)$?

Simplifications and Abbreviations

- Let us pick the last formula apart to understand it better (and also get a CNF representation along the way).
- Let us call the formula τ ("transition"):
  $\tau = o^i \rightarrow (v^i \leftrightarrow regr(v, eff(o))^{i-1})$.
- First, some abbreviations:
  - Let $e = eff(o)$.
  - Let $\rho = regr(v, e)$ ("regression").
    We have $\rho = trig(v, e) \vee (v \wedge \neg trig(\neg v, e))$.
  - Let $\alpha = trig(v, e)$ ("added").
  - Let $\delta = trig(\neg v, e)$ ("deleted").

$\rightsquigarrow$ $\tau = o^i \rightarrow (v^i \leftrightarrow \rho^{i-1})$ with $\rho = \alpha \vee (v \wedge \neg\delta)$

Picking it Apart (1)

Reminder: $\tau = o^i \rightarrow (v^i \leftrightarrow \rho^{i-1})$ with $\rho = \alpha \vee (v \wedge \neg\delta)$

$$\begin{aligned}\tau &= o^i \rightarrow (v^i \leftrightarrow \rho^{i-1})\\ &\equiv o^i \rightarrow ((v^i \rightarrow \rho^{i-1}) \wedge (\rho^{i-1} \rightarrow v^i))\\ &\equiv \underbrace{(o^i \rightarrow (v^i \rightarrow \rho^{i-1}))}_{\tau_1} \wedge \underbrace{(o^i \rightarrow (\rho^{i-1} \rightarrow v^i))}_{\tau_2}\end{aligned}$$

$\rightsquigarrow$ consider this two separate constraints τ_1 and τ_2

Picking it Apart (2)

Reminder: $\tau_1 = o^i \rightarrow (v^i \rightarrow \rho^{i-1})$ with $\rho = \alpha \vee (v \wedge \neg\delta)$

$$\begin{aligned}\tau_1 &= o^i \rightarrow (v^i \rightarrow \rho^{i-1})\\ &\equiv o^i \rightarrow (\neg\rho^{i-1} \rightarrow \neg v^i)\\ &\equiv (o^i \wedge \neg\rho^{i-1}) \rightarrow \neg v^i\\ &\equiv (o^i \wedge \neg(\alpha^{i-1} \vee (v^{i-1} \wedge \neg\delta^{i-1}))) \rightarrow \neg v^i\\ &\equiv (o^i \wedge (\neg\alpha^{i-1} \wedge (\neg v^{i-1} \vee \delta^{i-1}))) \rightarrow \neg v^i\\ &\equiv \underbrace{((o^i \wedge \neg\alpha^{i-1} \wedge \neg v^{i-1}) \rightarrow \neg v^i)}_{\tau_{11}} \wedge \underbrace{((o^i \wedge \neg\alpha^{i-1} \wedge \delta^{i-1}) \rightarrow \neg v^i)}_{\tau_{12}}\end{aligned}$$

$\rightsquigarrow$ consider this two separate constraints τ_{11} and τ_{12}

Interpreting the Constraints (1)

Can we give an intuitive description of τ_{11} and τ_{12}?
$\rightsquigarrow$ Yes!

- $\tau_{11} = (o^i \wedge \neg\alpha^{i-1} \wedge \neg v^{i-1}) \rightarrow \neg v^i$

  "When applying o, if v is false and o does not add it, it remains false."
  - called negative frame clause
  - in clause form: $\neg o^i \vee \alpha^{i-1} \vee v^{i-1} \vee \neg v^i$
- $\tau_{12} = (o^i \wedge \neg\alpha^{i-1} \wedge \delta^{i-1}) \rightarrow \neg v^i$

  "When applying o, if o deletes v and does not add it, it is false afterwards." (Note the add-after-delete semantics.)
  - called negative effect clause
  - in clause form: $\neg o^i \vee \alpha^{i-1} \vee \neg\delta^{i-1} \vee \neg v^i$

For STRIPS tasks, these are indeed clauses. (And in general?)

Picking it Apart (3)

Almost done!

Reminder: $\tau_2 = o^i \rightarrow (\rho^{i-1} \rightarrow v^i)$ with $\rho = \alpha \vee (v \wedge \neg\delta)$

$$\begin{aligned}\tau_2 &= o^i \rightarrow (\rho^{i-1} \rightarrow v^i)\\ &\equiv (o^i \wedge \rho^{i-1}) \rightarrow v^i\\ &\equiv (o^i \wedge (\alpha^{i-1} \vee (v^{i-1} \wedge \neg\delta^{i-1}))) \rightarrow v^i\\ &\equiv \underbrace{((o^i \wedge \alpha^{i-1}) \rightarrow v^i)}_{\tau_{21}} \wedge \underbrace{((o^i \wedge v^{i-1} \wedge \neg\delta^{i-1}) \rightarrow v^i)}_{\tau_{22}}\end{aligned}$$

$\rightsquigarrow$ consider this two separate constraints τ_{21} and τ_{22}

Interpreting the Constraints (2)

How about an intuitive description of τ_{21} and τ_{22}?

- $\tau_{21} = (o^i \wedge \alpha^{i-1}) \rightarrow v^i$

  "When applying o, if o adds v, it is true afterwards."
  - called positive effect clause
  - in clause form: $\neg o^i \vee \neg \alpha^{i-1} \vee v^i$
- $\tau_{22} = (o^i \wedge v^{i-1} \wedge \neg \delta^{i-1}) \rightarrow v^i$

  "When applying o, if v is true and o does not delete it, it remains true."
  - called positive frame clause
  - in clause form: $\neg o^i \vee \neg v^{i-1} \vee \delta^{i-1} \vee v^i$

For STRIPS tasks, these are indeed clauses. (But not in general.)

SAT Formula: Transitions

SAT Formula: Transitions

precondition clauses:

- $\neg o^i \vee pre(o)^{i-1}$ for all $1 \leq i \leq T$, $o \in O$

positive and negative effect clauses:

- $\neg o^i \vee \neg \alpha^{i-1} \vee v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$
- $\neg o^i \vee \alpha^{i-1} \vee \neg \delta^{i-1} \vee \neg v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$

positive and negative frame clauses:

- $\neg o^i \vee \neg v^{i-1} \vee \delta^{i-1} \vee v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$
- $\neg o^i \vee \alpha^{i-1} \vee v^{i-1} \vee \neg v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$

where $\alpha = trig(v, eff(o))$, $\delta = trig(\neg v, eff(o))$.

For STRIPS, all except the precondition clauses are in clause form.

The precondition clauses are easily convertible to CNF (one clause $\neg o^i \vee v^{i-1}$ for each precondition atom v of o).

C5.5 Summary

Summary: Sequential SAT Encoding (1)

Sequential SAT Encoding (1)

initial state clauses:

- v^0 for all $v \in V$ with $I(v) = \mathbf{T}$
- $\neg v^0$ for all $v \in V$ with $I(v) = \mathbf{F}$

goal clauses:

- γ^T

operator selection clauses:

- $o_1^i \vee \cdots \vee o_n^i$ for all $1 \leq i \leq T$

operator exclusion clauses:

- $\neg o_j^i \vee \neg o_k^i$ for all $1 \leq i \leq T$, $1 \leq j < k \leq n$

Summary: Sequential SAT Encoding (2)

Sequential SAT Encoding (2)

precondition clauses:

- $\neg o^i \vee pre(o)^{i-1}$ for all $1 \leq i \leq T$, $o \in O$

positive and negative effect clauses:

- $\neg o^i \vee \neg \alpha^{i-1} \vee v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$
- $\neg o^i \vee \alpha^{i-1} \vee \neg \delta^{i-1} \vee \neg v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$

positive and negative frame clauses:

- $\neg o^i \vee \neg v^{i-1} \vee \delta^{i-1} \vee v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$
- $\neg o^i \vee \alpha^{i-1} \vee v^{i-1} \vee \neg v^i$ for all $1 \leq i \leq T$, $o \in O$, $v \in V$

where $\alpha = trig(v, eff(o))$, $\delta = trig(\neg v, eff(o))$.

Summary

- SAT planning (planning as satisfiability) expresses a sequence of bounded-horizon planning tasks as SAT formulas.
- Plans can be extracted from satisfying assignments; unsolvable tasks are challenging for the algorithm.
- For each time step, there are propositions encoding which state variables are true and which operators are applied.
- We describe a basic sequential encoding where one operator is applied at every time step.
- The encoding produces a CNF formula for STRIPS tasks.
- The encoding follows naturally (with some work) from using regression to link state variables in adjacent time steps.